

BAYESIAN NONPARAMETRIC ESTIMATION OF MILKY WAY MODEL PARAMETERS USING A NEW MATRIX-VARIATE GAUSSIAN PROCESS BASED METHOD

BY DALIA CHAKRABARTY^{*,†,¶,||}, MUNMUN BISWAS^{‡,**}, SOURABH BHATTACHARYA^{§,††},
University of Warwick[¶] and University of Leicester^{||}, Indian statistical Institute^{**}, Indian Statistical Institute^{††}

The modelling of the Milky Way galaxy is an integral step in the study of galactic dynamics; this is because knowledge of model parameters that define the Milky Way directly influences our understanding of the evolution of our galaxy. Since the nature of the Galaxy's phase space in the neighbourhood of the Sun is affected by various Milky Way features, measurements of phase space coordinates of individual stars that live in this neighbourhood, will bear information about such features. Thus, inversion of such measurements can help us learn the parameters that describe such Milky Way features.

In the past this has been attempted via the "calibration method", a data and computational cost intensive method that is limited by logistical considerations. We argue in this paper that the calibration method is unsatisfactory because of reasons both methodological and computational. Here we develop a Bayesian inverse problem approach, where we model the available stellar velocity information matrix as an unknown function of the Milky Way model parameters, where this function is inverted using Bayesian techniques to predict the model parameters. This unknown function turns out to be matrix-variate, which we model as a matrix-variate Gaussian Process. We develop a general method to perform inverse, nonparametric learning, using such Gaussian Processes. For the inference we use the recently advanced Transformation-based Markov Chain Monte Carlo (TMCMC).

Application of our method to observed stellar velocity data results in estimates that are consistent with those in astrophysical literature. That we could obtain these results using far smaller data sets compared to those required for the calibration approach, is encouraging in terms of projected applications to external galaxies.

1. Introduction. Curiosity about the nature of the phase space that we earthlings live in, is only natural. On astronomical length scales, this translates to the pursuit of the topology of phase space in the Solar neighbourhood. By phase space of the stars in the Milky Way, is implied $\mathcal{W}$, the space of all states that the stars in our galaxy can attain. This reduces to the spatial $\mathbf{X}$ and velocity $\mathbf{P}$ vectors of all the stars in the Galaxy, where the spatial location of the a -th star is $\mathbf{X}_a = (X_1^{(a)}, X_2^{(a)}, X_3^{(a)})^T$, and its velocity vector $\mathbf{P}_a = (P_1^{(a)}, P_2^{(a)}, P_3^{(a)})^T$.

The phase space of the Milky Way in the vicinity of the Sun, has been studied and astrophysical modelling indicates that in this local patch, the phase space structure is sculpted by different dynamical

^{*} Associate Research fellow at Department of Statistics, University of Warwick

[†] Lecturer of Statistics at Department of Mathematics, University of Leicester

[‡] PhD student in Statistics and Mathematics Unit, Indian Statistical Institute

[§] Assistant Professor in Bayesian and Interdisciplinary Research Unit, Indian Statistical Institute

Keywords and phrases: Supervised learning,, Inverse problems,, Gaussian Process,, Matrix-variate distributions,, Transformation-based MCMC

features of the Milky Way (Antoja et al., 2009; Chakrabarty, 2007; Minchev et al., 2009), such as a central, rotating, stellar bar in the Galaxy as well as the spiral pattern of the Galaxy. Thus, the phase portrait of the solar neighbourhood is affected by the parameters that define these Galactic features. Relevant parameters include the orientation of the observer at Earth (i.e. approximately at the Sun) with respect to an axis of the bar, the distance from us (i.e. the Sun) to the centre of the Galaxy, structural parameters of the bar and the spiral pattern, as well as parameters that describe the dynamics of these features, such as the frequencies of rotation of the spiral pattern Ω_s and that of the bar Ω_b , etc. In fact, there are further features that affect the Galactic phase space structure so that the feature parameter space demands expansion to include parameters that have hitherto been ignored from the modelling of the Galactic phase space.

Given that these feature parameters affect the local patch of our Galaxy, if phase space coordinates of a sample of stars that live in the local patch are measured, such data will bear information about the handiwork of these relevant features. It then follows that inversion of such phase space information will allow for the learning of the unknown feature parameters. This approach has been adopted in the modelling of our galaxy, to result in the estimation of the angular separation of the Sun from a chosen axis of the bar and the distance of the Sun from the Galactic centre (Dehnen, 2000; Fux, 2001; Minchev et al., 2010; Simone, Wu and Tremaine, 2004). The other relevant feature parameters are typically held constant in such modelling.

The modelling of the solar neighbourhood by Chakrabarty (2007) is similar in that the two-dimensional solar location is estimated. The method used in this work is an example of the “calibration method”. The objective in this work was to predict the solar location given the available stellar velocity measurements and simulated phase space data of a sample of stars generated at n different chosen values of the Milky Way feature parameter vector $\mathbf{S}$. Each of the n simulated velocity data was employed to generate n density estimates in the space of the velocity vectors, via kernel density estimation techniques. The estimated density functions appeared highly multi-modal and heterogeneous. A non-parametric frequentist test was designed to test for the null that the observed data is distributed according to the i -th density function that is estimated from the i -th simulated data set that was generated at the i -th chosen value of the Milky Way model parameter vector $\mathbf{S}$, ($i = 1, \dots, n$), (Chakrabarty, 2011). The p -value of the used test statistic was recorded for each i . Those choices of the value of $\mathbf{S}$ that corresponded to high p -values, were considered better supported by the observed data. Hence the empirical distribution of these p -values in the space of $\mathbf{S}$, obtained from the n repeated testing, was used to provide interval estimates of the Milky Way parameter vector.

Since the best match is sought over a very large collection of training data points - obtained by constructing a grid over the relevant sample space - the method requires computational effort and resources. This shortcoming had compelled Chakrabarty (2007) to resort to an unsatisfactory coarse gridding of the space of $\mathbf{S}$. This problem gets acute enough for the method to be rendered useless when the dimensionality of the vector $\mathbf{S}$ that we hope to learn, increases. Moreover, the method of quantification of uncertainty of the estimate of the location is also unsatisfactory, dependent crucially on the binning details of the space of $\mathbf{S}$, which in turn is bounded by cost and memory considerations.

Thus, an improved method of learning the unknown Milky Way model parameter vector is sought. In this paper, we present such a method that allows for the Bayesian non-parametric learning of an unknown model parameter vector, given matrix-variate data. Such data is then expressed as a matrix-variate function of the model parameter vector where this unknown function is modelled as a realisation of a Gaussian Process that we discuss here. We demonstrate the effectiveness of this method

even with much smaller data sets than were used in astronomical modelling in the past (Chakrabarty, 2007). That the new method is able to work with smaller phase space data sets, is an important benefit, particularly in extending the application to galaxies other than our own, in which small numbers of individual stars are going to be tracked in the very near future for their velocities, under observational programmes such as GAIA (Kucinskas, Lindegren and Vansevičius, 2005; Lindegren et al., 2007, <http://www.rssd.esa.int/index.php?project=GAIA&page=index>) and PANStarrs (Johnston, Bullock and Strauss, 2009); the sample sizes of measured stellar velocity vectors in these programmes will be much smaller in external galaxies than what has been possible in our own. The other major advantage of this presented method is that it readily allows for the expansion of dimensionality of the feature parameter vector of the studied galaxy and is capable of taking errors in the measurements into account, within a Bayesian approach.

Of course, it is of aesthetic interest to learn our location inside our galaxy, but the relevance of this exercise goes much beyond. Any modelling of the Milky Way invokes the solar location - in that sense, the solar location is of vital importance in the study of the Milky Way. Thus, learning the distance of the Sun from the galactic centre, $R_{\odot}$, will crucially constrain the rotational frequency Ω_b of the bar in the Milky Way, as we discuss later in Section 5.4. The value of Ω_b affects the understanding of some crucial physical processes in the Milky Way. Also, the direction $\Phi = 0$ is chosen in our models to coincide with the long axis of the central stellar bar in the Galaxy. Thus, learning the angular separation of the Sun from this chosen axis, $\Phi_{\odot}$, would allow for an understanding of the separation of the line joining the Sun to the Galactic centre, from the long axis of the bar; this in turn will allow for improved understanding of the effect of the bar on the dynamics of the solar neighbourhood (Englmaier and Gerhard, 1999; Fux, 2001; Minchev et al., 2010).

The rest of the paper is structured as follows. In Section 2 we discuss the structure of the available simulated and real data. The salient aspects of our application are delineated in Section 3 which also includes the motivation for a Gaussian Process-based estimation. In Section 4 we present the details of the modelling strategy that we adopt. In this context, the relationship between the information and the model parameter is presented as a realisation of a Gaussian Process that we discuss. The achievement of the posterior probability density of the unknowns, given all data, is arrived at in Section 4.4, following discussions about the likelihood and priors on the unknown parameters in Section 4.1 and Section 8 respectively. The treatment of measurement errors within the modelling is discussed in Section 4.5. Section 5 deals with the implementation of the method to solve the real-life astronomical problem. In particular, details of the inference strategem used in our work are given in Section 5.1, Section 5.2 and Section 5.3. and Section 5.4 contains results obtained from using the available data. We compare the obtained results with the estimates available in the astronomical literature in Section 5.5 and focus on the very useful advantage of our methodology (that it can work even with much smaller data sets than other available) in Section 5.6. The benefits of our method are delineated subsequently in Section 5.7. The paper is rounded up with a discussions section which includes a discourse on other possible areas of application of our method.

2. Data. Essentially, we are attempting an inverse problem, namely that of estimating the unknown Milky Way model parameter vector $\mathbf{S}$ from the sampled discrete phase space stellar data. The method is designed to work with observed real data as well as simulated data, given that in addition to the observed data, n simulated stellar phase space data sets are available (Chakrabarty, 2007) at n chosen values of $\mathbf{S}$, for 4 distinct astrophysical models of the Galaxy. The method is designed to estimate the Galactic parameters $\mathbf{S} \in \mathbb{R}^d$, that affect the structure of the phase space around the Sun. In our application, we

restrict the dimensionality of $\mathcal{S}$ to 2, i.e. we estimate the coordinates of the radial location $R_\odot$ of the Sun and the solar azimuthal coordinate $\Phi_\odot$.

It is to be noted that we approximate the geometry of the Milky Way disc as a 2-dimensional disc. Thus, we confine analysis to such a two-dimensional spatial geometry, rendering the spatial vector of the a -th star $(X_1^{(a)}, X_2^{(a)})^T$, and the velocity vector $(P_1^{(a)}, P_2^{(a)})^T$. Given the disk spatial geometry, we find it sometimes useful to use the polar, rather than the Cartesian coordinate system to scan the Galaxy. Then the radial coordinate of a particle is R and azimuthal coordinate is Φ , such that $X_1 = R \cos \Phi$, $X_2 = R \sin \Phi$, $R \geq 0$, $\Phi \in [0, 360^\circ]$. The radial and transverse components of the velocity vector are then U and V .

While the Milky Way model parameters are sought with respect to the centre of the Galaxy, the measurements that are available, have been taken by us, i.e. with respect to the Sun. In other words, the measured stellar velocities $(U^{(a)}, V^{(a)})^T$, $a = 1, \dots, j$, are with respect to the solar velocity while the measured spatial locations of the stars are with respect to the location of the Sun. Furthermore, the spatial locations of the sampled stars are within a small distance ϵ from the Sun (Fux, 2001). We interpret these data to have been sampled from a circle with radius ϵ centred at the Sun, living in the space of the Milky Way disk. Also, the angular distribution of the sampled stars, inside this 2-ball, is approximately uniform. Thus, the summary of the distribution of the measured spatial locations of the sampled stars will always coincide with the origin of the used spatial coordinate system, irrespective of what the galactocentric spatial location of this origin is, i.e. irrespective of the observer's galactocentric spatial location. Thus, these measured spatial locations cannot constrain the sought galactocentric location of the Sun.

So the measurements (from the Sun) of spatial locations of the sampled stars do not bear any information about the unknown $(R_\odot, \Phi_\odot)^T$, though the measured heliocentric stellar velocities are indeed affected by the choice of $(R_\odot, \Phi_\odot)^T$. This is because, a given star, if observed from different spatial locations in the Galaxy, would appear to move in different ways. For example, if a star appears to have a velocity vector directed along the line that joins itself to the observer at point A on the Milky Way disc, this observer will register its velocity to be entirely radial, with zero angular component of the velocity vector. On the contrary, had the observer been at a different point B , the velocity vector of this star would have registered to have had a radial as well as an angular component, in general. Thus, the observed stellar velocities will bear information about the location of the observer, i.e. the Sun. Then the available velocity information $\mathbf{V}$ can be considered to bear the signature of Milky Way parameters $\mathcal{S}$. In principle, beyond just the galactocentric solar location, if there are Milky Way parameters that physically affect the structure of the phase space, velocity information drawn from that phase space will bear the signature of such Milky Way parameters.

In light of the above discussion, the measured data used in our work is considered to comprise stellar velocity information, $\{(U_a, V_a)^T\}_{a=1}^j$, where a total of j number of stars are observed. Thus, the measured velocity information is a matrix but in our modelling, we find it more useful to consider the j number of measurements of the k -dimensional velocity vectors, as a jk -dimensional velocity vector $\mathbf{v}^{(test)}$. In our application to learn the Milky Way model parameters, the dimensionality of the measured velocity vectors is 2, i.e. $k=2$.

As mentioned above, we supplement information about the unknown Milky Way parameters with simulated information that is generated from dynamical simulations of 4 astronomical models of our Galaxy, at n distinct choices of values of $\mathcal{S}$. In fact, the simulated data generated for the q -th astronomical model, corresponding to chosen values, $s_1^*, \dots, s_n^*$ of $\mathcal{S}$, comprise the training data $\mathcal{D}_s^{(q)}$,

$q = 1, 2, 3, 4$. At the i -th chosen value, $\mathbf{s}_i^*$ of $\mathbf{S}$, let there be j number of k -dimensional velocity vectors simulated, $i = 1, \dots, n$. Thus, we will expect such simulated information to be in the form of a series of n matrices, each of which is $j \times k$ -dimensional. However, as with the real data, we find it useful in our modelling to treat each simulated velocity matrix as a jk -dimensional vector, and there are n such generated velocity vectors, generated at the n different choices of the model parameter vector $\mathbf{S}$. From this viewpoint, the entire simulated velocity information is a matrix $\mathcal{D}_s$, with n rows and jk columns, i.e. the simulated velocity information can be considered to be the matrix $\mathcal{D}_s^{(n \times jk)}$, such that the i -th row is the jk -dimensional simulated velocity vector $\mathbf{v}_i$, generated at the i -th chosen value $\mathbf{s}_i^*$ of $\mathbf{S}$, where $\mathbf{S}$ is itself a d -dimensional vector.

In our work, the simulated $\mathcal{D}_s$ is referred to as the training data while the real or observed data $\mathbf{v}^{(test)}$ is referred to as test data. This is because the simulated data are generated at known values of $\mathbf{S}$ while the model parameter supported by the real data is unknown. The set $\{\mathbf{s}_1^*, \dots, \mathbf{s}_n^*\}$ of the n chosen values of $\mathbf{S}$ at which the simulated velocities are generated, is referred to as the design set and $\mathbf{s}_i$ as the i -th design vector.

The inverse method that we advance below is a generic methodology that can be applied to estimate the d -dimensional vector $\mathbf{S}$, where d is bounded from below by the number of parameters in the astrophysical models that can be called upon to bear influence on the phase space structure of the Galaxy in the solar neighbourhood. As we have said above, we demonstrate an application of our developed learning technique to the learning of the 2-dimensional solar location vector. The reason for restricting our application to the case of $d=2$ is the existence of stellar phase space data generated in dynamical simulations of astrophysical models of the Milky Way which involve scanning over chosen guesses for $R_\odot$ and $\Phi_\odot$, with all other feature parameters held constant. If simulated data distinguished by choices of other Milky Way model feature parameters become available, then the implementation of such data as training data will be possible, allowing then for the learning of Milky way model parameters in addition to $R_\odot$ and $\Phi_\odot$. The methodology that we advance is indeed a generic one, with computational costs being the only concern in extending to cases of $d > 2$; that extending to a higher dimensional $\mathbf{S}$ only linearly scales computational costs, is discussed later in Section 6.

Actually, the learning of the unknown $\mathbf{S}$ is attempted 4 times, each time using the observed data $\mathbf{v}^{(test)}$ and one of the 4 distinct $\mathcal{D}_s^{(q)}$ data that are generated at the chosen design set, using the 4 astronomical models that were considered by Chakrabarty (2007), $q = 1, 2, 3, 4$. Above, the training data $\mathcal{D}_s^{(q)}$ is the $n \times jk$ -dimensional matrix that is simulated using the q -th astrophysical model of the Milky Way, such that its i -th row is the i -th velocity vector $\mathbf{v}_i$ that is generated at the choice of $\mathbf{S} = \mathbf{s}_i^*$ $i = 1, \dots, n$. The choice of the base astrophysical model is distinguished by the ratio of the rotational frequencies of the spiral to the bar, Ω_s/Ω_b . That this ratio is relevant to the phase space structure of the model of the Galaxy is due to the fact that Ω_s/Ω_b can crucially control the degree of chaos in the Galactic model. For example, it is well known in chaos theory that when Ω_s/Ω_b is such that one of the radii at which the bar and the stellar disk resonate, concurs with a radius at which the spiral and the stellar disk resonate, global chaos is set up in the system (G. Walker and J. Ford, 1969). Chakrabarty and Sideris (2008) have corroborated that the degree of chaos is maximal in the astrophysical Galactic model marked by such a ratio ($\Omega_s/\Omega_b=22/55$). They report that in models marked by slightly lower ($\Omega_s/\Omega_b=18/55$) or higher ($\Omega_s/\Omega_b = 25/55$) values of this ratio, chaos is still substantial. In the Galactic model that precludes the spiral however, chaos was quantified to be minimal. It is these 4 states of chaos - driven by the 4 values of Ω_s/Ω_b - that mark the 4 astrophysical models as distinct. Thus, the 4 astrophysical models lead to model phase space structures that are differently chaotic. This results in 4 distinct simulated

velocity information sets $\mathcal{D}_s^{(1)}, \mathcal{D}_s^{(2)}, \mathcal{D}_s^{(3)}, \mathcal{D}_s^{(4)}$ that bear the effects of such varying degrees of chaos, each generated at the chosen design set $\{s_1^*, \dots, s_n^*\}$. As attempts to learn the unknown Galactic model parameter vector are undertaken with one astrophysical base model at a time, each such attempt is independent of the other. Consequently, we simplify the notation in our discussion of the method below, by dropping the reference to the index q on the training data sets hereafter.

Repeated learning of the unknown $s^{(new)}$ using the real measurements $\mathbf{v}^{(test)}$ and one of these simulated data sets will allow for the examination of the influence of a chaotic phase space structure on such learning. For instance, if the implementation of simulated data sets that are generated from base astrophysical models marked by similar chaoticities, result in learnt values of $s^{(new)}$ that are widely disparate, that would be cause for concern. At the same time, a method that is unable to perform successful learning of $s^{(new)}$ in the face of extreme chaos - such as the method of Chakrabarty (2007) - is viewed as weak. We show below that in our method, the robust model uncertainty estimation within a Bayesian framework allows for the learning even in the face of global chaos in the base astrophysical model. Details of the dynamical simulations performed on the 4 astrophysical models are given in the supplementary section S-1.

3. Case study. In this work, our interest lies in predicting the value of the unknown Milky Way model parameter $s^{(new)}$ that supports the real data vector $\mathbf{v}^{(test)}$, given the training data matrix $\mathcal{D}_s$ that is generated at chosen values $s_1^*, \dots, s_n^*$, of the Milky Way model parameter vector $\mathbf{S}$. We define the chosen design set as $\{s_1^*, \dots, s_n^*\}$. The training data is employed to train the model for the unknown functional relationship between the available velocity information and the set of values of the model parameter vector that support this information. We present the learning of the solar position in the Milky Way disk by inverting real stellar velocity measurements in conjunction with simulated stellar velocity data generated at a chosen design set. Then we are interested in the inverse problem of predicting $s^{(new)}$ given $\mathcal{D}_s, \mathbf{v}^{(test)}$ and the trained model for the aforementioned functional relationship.

For this inversion, one possible way is to first estimate this function from the data, using well-known nonparametric techniques, and then use numerical methods to achieve the inverse solution. The complexity of both these exercises - particularly the latter - increases as the dimensionality of the function space increases, unless simplifying assumptions are made. In fact, modelling high-dimensional functions using splines/wavelets require restrictive assumptions and in particular, may fail to adequately take into account the correlation structure between the component functions. Furthermore, this approach has been criticised on the ground that it ignores parameter uncertainty, in that the approach provides only a point estimate of the set of model parameters.

A Bayesian equivalent of this approach could involve the modelling of the unknown underlying function with splines or wavelets, and then computing the posterior distribution of $s^{(new)}$, conditional on the training and the test data sets, after marginalising over the other model parameters. This computation, as well as the marginalisation, are in principle straightforward to implement using well-known Markov Chain Monte Carlo (MCMC) methods. Thus, the Bayesian approach avoids the difficulties of nonparametric model parameter estimation and subsequent inversion. Additionally, the Bayesian approach does not ignore parameter uncertainty while predicting the unknown location. Furthermore, priors on the unknown parameters can bring in extra information into the model. Thus, in principle, a training data set of even smaller size than that required in the classical approach, will be adequate in the Bayesian paradigm. However, as already discussed above, modelling high-dimensional functions using splines/wavelets is somewhat restrictive in that this approach fails to provide an appropriate correlation structure for the component-wise functions.

In this paper, we circumvent this problem by introducing a Gaussian Process approach to modelling the high-dimensional functional relationship between available information vectors $\mathbf{V}$ and the value of $\mathbf{S}$ that supports this information. The velocity information that results from collating such information vectors is matrix-variate. The functional relationship between a general information matrix $\mathcal{V}$ and the corresponding matrix of values of $\mathbf{S}$ is modelled in our work with a Gaussian Process. In the face of the matrix-variate nature of the information, this functional relationship needs to be modelled as a realisation from a matrix-variate Gaussian Process ($\mathcal{GP}$).

Indeed, had we been treating the information $\mathbf{V}$ generated at a value of $\mathbf{S}$ as a $j \times k$ -dimensional matrix, we would have modelled the functional relation between $\mathbf{S}$ and $\mathbf{V}$ by a $j \times k$ -dimensional matrix-variate $\mathcal{GP}$. In that case, one realisation from such a matrix-variate $\mathcal{GP}$ would have been the n number of $j \times k$ -dimensional matrices containing velocity values that are generated at n values of $\mathbf{S}$. However, as said before, in our treatment, the j number of k -dimensional velocity vectors generated at any value of $\mathbf{S}$ is treated as a jk -dimensional vector. In this paradigm, each of the n such information vectors that will be considered to be a realisation from a vector-variate $\mathcal{GP}$ of dimensionality jk . Thus, our view allows for the reduction of the problem from a matrix-variate $\mathcal{GP}$ to a vector-variate $\mathcal{GP}$.

So we treat the general $j \times k$ -dimensional stellar velocity information as a jk -dimensional information vector $\mathbf{V}^{(jk \times 1)}$. Then we model $\mathbf{V}$ as a function of the value s of a general model parameter vector $\mathbf{S}$ at which such information is generated. Thus, we write $\mathbf{V} = \xi(s)$ where s is the value of $\mathbf{S}$ that supports this information vector $\mathbf{V}$. To be precise, in our application, each stellar velocity vector is 2-dimensional, i.e. $k=2$. Also, we seek to learn a model parameter vector that is 2-dimensional, i.e. $d=2$. In the Bayesian approach that we adopt, a much smaller j ($=50$) allows for inference on the unknown value $s^{(new)}$ of the Milky Way model parameter vector, than $j \sim 3000$ that is demanded by the aforementioned calibration approach used by Chakrabarty (2007). The simulated data presented in Chakrabarty (2007) that we use here, is generated at 216 distinct values of $\mathbf{S}$, i.e. $n=216$.

Thus, our design set comprises 216 chosen values $s_1^*, \dots, s_n^*$ of the model parameter vector. For each of the 4 base astrophysical models, at each chosen s_i^* , 50 2-dimensional stellar velocity vectors are randomly drawn from the velocity data that are generated from dynamical simulations performed at that value of $\mathbf{S}$. Here, $i = 1, \dots, 216$. As explained before, the 216 pieces of velocity information generated in this way are treated in our work as vectors $\mathbf{v}_1, \dots, \mathbf{v}_{216}$, where $\mathbf{v}_i$ has dimensionality $50 \times 2 = 100$ for each i . Then the training data in our work comprises all such vectors and is represented as $\mathcal{D}_s^{(216 \times 100)}$. The real data is treated in our work as the 100-dimensional vector $\mathbf{v}^{(test)}$. The i -th design vector is s_i^* which is a d -dimensional vector $\forall i = 1, \dots, n$.

It will be seen in Section 5.4 that, in spite of our relatively small number of stars in the data sets, the summary of the posterior distributions of the unknown solar position is consistent with existing knowledge in the astrophysics literature. The other very important contribution of our work is that it allows for the learning of a high-dimensional model parameter vector.

4. Modelling. In line with the discussion in Neal (1998), in our application we have n jk -dimensional simulated velocity vectors (inputs), each generated at a chosen value of the model parameter vector (target), i.e. we have the n observations $(\mathbf{v}_1, s_1^*), \dots, (\mathbf{v}_n, s_n^*)$, (where s_i^* is a known d -dimensional vector, $\forall i = 1, \dots, n$). Our aim is to predict the unknown Milky Way model parameter vector $s^{(new)}$ when we have only information about the input in the form of the real velocity vector $\mathbf{v}^{(test)}$. A predictive distribution of $s^{(new)}$ is sought, conditioned on the real velocity vector and the training data generated at the known design vector values, namely at $s_1^*, \dots, s_n^*$. This is achieved in an approach in which we treat

the input - that is to begin with matrix-variate - alternatively as a vector of corresponding dimensions, allowing us to then consider the functional relation between the inputs and the targets as a realisation from a vector-variate $\mathcal{GP}$. In this section we ignore measurement errors and present our model of these n vector-variate functions. Later in Section 4.5, we delineate the method used to take measurement uncertainties on board.

We define the full information matrix $\mathcal{V}^{(n \times jk)}$ as one with the i -th row given by the velocity information vector $(\mathbf{V}_i^{(jk \times 1)})^T$, where $i = 1, \dots, n$. Then $\mathbf{V}_i$ is supported by the value s_i of the model parameter vector $\mathbf{S}$, so that the equation $\mathbf{V} = \boldsymbol{\xi}(\mathbf{s})$ expresses the functional relationship between the two variables $\mathbf{V}$ and $\mathbf{S}$. We express the $i\ell$ -th element of the information matrix $\mathcal{V}$ as $V_{i\ell}$ which is equal to the ℓ -th component function $\xi_\ell(s_i)$ of the jk -dimensional $\boldsymbol{\xi}(s_i)$. Then $\xi_\ell(s_i)$ is a scalar valued function of the vector s_i , where $i = 1, \dots, n$ and $\ell = 1, \dots, jk$. In other words,

$$\begin{pmatrix} V_{11} & V_{12} & \dots & V_{1jk} \\ V_{21} & V_{22} & \dots & V_{2jk} \\ \dots & \dots & \dots & \dots \\ V_{n1} & V_{n2} & \dots & V_{njk} \end{pmatrix} = \begin{pmatrix} \xi_1(s_1) & \xi_2(s_1) & \dots & \xi_{jk}(s_1) \\ \xi_1(s_2) & \xi_2(s_2) & \dots & \xi_{jk}(s_2) \\ \dots & \dots & \dots & \dots \\ \xi_1(s_n) & \xi_2(s_n) & \dots & \xi_{jk}(s_n) \end{pmatrix}$$

Thus, the i -th row of the $\mathcal{V}^{(n \times jk)}$ matrix is the jk -dimensional vector $\mathbf{V}_i$ that is set equal to $\boldsymbol{\xi}(s_i) := (\xi_1(s_i), \dots, \xi_{jk}(s_i))^T$, $\forall i = 1, \dots, n$. We model the jk -dimensional function $\boldsymbol{\xi}(\cdot)$ as a jk -dimensional vector-variate $\mathcal{GP}$. One realisation from this $\mathcal{GP}$ is $\{\boldsymbol{\xi}(s_1), \boldsymbol{\xi}(s_2), \dots, \boldsymbol{\xi}(s_n)\}$. Then the joint distribution of $\{\boldsymbol{\xi}(s_1), \boldsymbol{\xi}(s_2), \dots, \boldsymbol{\xi}(s_n)\}$ is matrix normal, with adequate parametrisation. We represent this as

$$(4.1) \quad \{\boldsymbol{\xi}(s_1), \boldsymbol{\xi}(s_2), \dots, \boldsymbol{\xi}(s_n)\} \sim \mathcal{MN}_{n,jk}(\boldsymbol{\mu}, \mathbf{A}, \boldsymbol{\Omega}),$$

where the mean matrix of this matrix normal distribution is the $n \times jk$ -dimensional matrix $\boldsymbol{\mu}$, the left covariance matrix is the $n \times n$ -dimensional $\mathbf{A}$ and the right covariance matrix is the $jk \times jk$ -dimensional matrix $\boldsymbol{\Omega}$. Before describing these individual matrix-variate parameters of this distribution below, we note that Equation 4.1 is the same as saying that the likelihood is matrix normal.

In our work we choose to implement the popularly used square exponential covariance function (Rasmussen and Williams, 2006; Santner, Williams and Notz, 2003; Scholkopf and Smola, 2002). This covariance function is easy to implement. It renders the sampled functions smooth and infinitely differentiable while the covariance structure itself is dependent only on the Euclidean distance between two values of the function variable so that the covariance is invariant to rotations and translations in the space of this variable. It is hard to preempt if such a stationary and isotropic covariance function is too restrictive to properly model correlations for the application at hand. While acknowledging such a possibility, we resort to relaxing the choice of a zero mean function though that is a popular choice. Instead we choose to define the mean function in a way that is equivalent to the suggestion that the velocity information is viewed as centred around a linear model with the residuals characterised by a vector-variate $\mathcal{GP}$ (A. O'Hagan, 1978; Cressie, 1993). We then integrate over all such possible global intercepts to arrive at a result that is more general than if the mean is fixed at zero. An advantage of the non-zero mean function is that in the limit of the smoothness parameters (characterising the smoothness of the functions sampled from this $\mathcal{GP}$) approaching large values, the random function reduces to a linear regression model, which is plausible. In models with a zero mean function though, in this limit of

very large smoothness, the random function will concur with the errors, which appears less intuitively pleasing.

The non-zero mean function of the $\mathcal{GP}$ is defined to be $\mu(\cdot) := \mathbf{H}\mathbf{B}$, where

$$(4.2) \quad \begin{aligned} \mathbf{H}^T &:= [\mathbf{h}^{(m \times 1)}(\mathbf{s}_1), \dots, \mathbf{h}^{(m \times 1)}(\mathbf{s}_n)], \quad \text{with} \\ \mathbf{h}^{(m \times 1)}(\cdot) &= (h_1(\cdot), \dots, h_m(\cdot))^T \end{aligned}$$

For $e = 1, \dots, m$, $h_e(\cdot)$ could be any function. However, in accordance to the suggestion that such a non-zero mean function be expressed in terms of a few basis functions (Rasmussen and Williams, 2006), we choose to fix this functional form as $h(\cdot) := (1, \cdot)^T$ for all values of $\mathbf{S}$. (A similar construct was used by Blight and Ott (1975) who performed a $\mathcal{GP}$ -based polynomial regression analysis). We also choose to set $m = d + 1$. Thus, in our treatment, $\mathbf{h}(\cdot)$ is a $(d + 1)$ -dimensional matrix. The extent to which this chosen mean function deviates from being zero is borne by the coefficient matrix $\mathbf{B}$ which is the other factor in the definition of the mean matrix. We define $\mathbf{B}$ as

$$(4.3) \quad \mathbf{B} = (\beta_{11}, \dots, \beta_{j1}, \dots, \beta_{1k}, \dots, \beta_{jk})$$

where for $p = 1, \dots, j$, $p' = 1, \dots, k$, $\beta_{pp'}$ is an m -dimensional column vector. As we choose to set $m = d + 1$, $\mathbf{B}$ is a matrix with $d + 1$ rows and jk columns.

The amplitude of the covariance function is characterised by the matrix $\mathbf{\Omega}$ which is $jk \times jk$ -dimensional and is defined as

$$(4.4) \quad \mathbf{\Omega} = \mathbf{\Sigma} \otimes \mathbf{C}$$

where $\mathbf{\Sigma}$ is the $k \times k$ covariance matrix for a given star in a row of the matrix-variate function $\xi(\cdot)$. Thus $\mathbf{\Sigma}$ corresponds to the covariance amongst components of the velocity vector of each star for a given value of $\mathbf{S}$. On the other hand, $\mathbf{C}$ is the $j \times j$ covariance matrix for a given component of the velocity vector for the stars at a fixed value of $\mathbf{S}$.

The left covariance matrix of the matrix-normal distribution defined in Equation 4.1 is $\mathbf{A}$ which is $n \times n$ -dimensional. Given the choice of square exponential covariance function, it is

$$(4.5) \quad \begin{aligned} \mathbf{A}^{(n \times n)} &:= [a(\cdot, \cdot)], \quad \text{where} \\ a(\mathbf{s}, \mathbf{s}') &\equiv \exp\{-(\mathbf{s} - \mathbf{s}')^T \mathbf{Q} (\mathbf{s} - \mathbf{s}')\}, \end{aligned}$$

for any 2 values $\mathbf{s}$ and $\mathbf{s}'$ of $\mathbf{S}$. Here, $\mathbf{Q}^{(d \times d)}$ represents the inverse of the scale length that underlies correlation between functions at any two values of the function variable. In other words, $\mathbf{Q}$ is the matrix of the smoothness parameters. Thus, $\mathbf{Q}$ is a matrix that bears information about the smoothness of the sampled functions; it is a diagonal matrix consisting of d non-negative smoothness parameters denoted by $b_1, \dots, b_d$. In other words, we assume the same smoothness for each component function of $\xi(\cdot)$. This smoothness is determined by the parameters $b_1, \dots, b_d$. We will learn these smoothness parameters in our work from the data. Of course, though we say that the smoothness is learnt in the data, the underlying effect of the choice of the square exponential covariance function on the smoothness of the sampled functions is acknowledged. Indeed, as Snelson (2007) states, one concern about the square exponential function is that it renders the functions sampled from it as artificially smooth. An alternative covariance function, such as the Matern class of covariances (Matern, 1986; Snelson, 2007; Tilmann Gneiting and William Kleiber and Martin Schlather, 2010), could give rise to sampled

functions that are much rougher than those obtained using the square exponential covariance function, for the same values of the hyperparameters of amplitude and scale that characterise these covariance functions (see Chapter 1 of Snelson's thesis).

Let $\omega_{r\ell}$ denote the (r, ℓ) -th element of Ω , $c_{r\ell}$ the (r, ℓ) -th element of C and let $\sigma_{r\ell}$ denote the (r, ℓ) -th element of Σ . Let the ℓ -th component function of $\xi(\cdot)$ be $\xi_\ell(\cdot)$ with $\ell = m_1 k + m_2$, where $\ell = 1, \dots, jk$ and $m_2 = 1, 2, \dots, k, m_1 = 0, 1, \dots, j-1$. Then the correlation between the components of $\xi(\cdot)$ yields the following correlation structures:

$$(4.6) \quad \text{corr}(\xi_{m_1 k + m_2}(\mathbf{s}_i), \xi_{m'_1 k + m_2}(\mathbf{s}_i)) = \frac{\sigma_{m_1 m'_1}}{\sqrt{\sigma_{m_1 m_1} \sigma_{m'_1 m'_1}}} \forall m_2, i \text{ and } m_1 \neq m'_1$$

$$(4.7) \quad \text{corr}(\xi_{m_1 k + m_2}(\mathbf{s}_i), \xi_{m_1 k + m'_2}(\mathbf{s}_i)) = \frac{c_{m'_2 m'_2}}{\sqrt{c_{m_2 m_2} c_{m'_2 m'_2}}} \forall m_1, i \text{ and } m_2 \neq m'_2$$

$$(4.8) \quad \text{corr}(\xi_{m_1 k + m_2}(\mathbf{s}_i), \xi_{m_1 k + m_2}(\mathbf{s}_i)) = \frac{c_{m_2 m'_2} \sigma_{m_1 m'_1}}{\sqrt{c_{m_2 m_2} \sigma_{m_1 m_1} c_{m'_2 m'_2} \sigma_{m'_1 m'_1}}} \forall i, m_1 \neq m'_1, m_2 \neq m'_2$$

$$(4.9) \quad \text{corr}(\xi_\ell(\mathbf{s}_1), \xi_\ell(\mathbf{s}_2)) = a(\mathbf{s}_1, \mathbf{s}_2) \forall \ell \text{ and } \mathbf{s}_1 \neq \mathbf{s}_2$$

4.1. Likelihood. Along these lines, the i -th row of the training data $\mathcal{D}_s$ that is itself $n \times jk$ -dimensional, can be expressed as a function of the i -th chosen design vector $\mathbf{s}_i^*$ of the Milky Way model parameter vector $\mathbf{S}$. Here the i -th row of $\mathcal{D}_s$ is the jk -dimensional vector $\mathbf{v}_i^T$ that is simulated at $\mathbf{S} = \mathbf{s}_i^*$, $i = 1, \dots, n$. Thus, we can write $\mathbf{v}_i = \xi(\mathbf{s}_i^*)$. Our ultimate aim is to learn the unknown value of $\mathbf{S}$ at which the real data $\mathbf{v}^{(test)}$ is generated. In this attempt, we write the probability distribution of the training data $\mathcal{D}_s$, given the parameters that define the distribution of $\xi(\cdot)$. In order to achieve this likelihood, we define

- the $n \times jk$ -dimensional mean function $\mathbf{H}_D \mathbf{B}$, where the linear form of the mean structure is contained in $\mathbf{H}_D^{(n \times m)} := [\mathbf{h}^{(m \times 1)}(\mathbf{s}_1^*), \dots, \mathbf{h}^{(m \times 1)}(\mathbf{s}_n^*)]$ and the coefficient matrix $\mathbf{B}$ is defined in Equation 4.3.
- the square exponential factor in the covariance matrix $\mathbf{A}_D^{(n \times n)} := [\exp\{-(\mathbf{s}^* - \mathbf{s}'^*)^T \mathbf{Q}(\mathbf{s}^* - \mathbf{s}'^*)\}]$ (see Equation 4.5).
- the amplitude of the covariance matrix $\Omega^{(jk \times jk)} = \Sigma^{(k \times k)} \otimes C^{(j \times j)}$ (see Equation 4.4),

Then it follows from the matrix normal distribution of in Equation 4.1, with mean function defined in Equation 4.2 and Equation 4.3, and covariance matrix defined using Equation 4.5 and Equation 4.4, that $\mathcal{D}_s$ is distributed as matrix normal with mean matrix $\mathbf{H}_D \mathbf{B}$, left covariance matrix $\mathbf{A}_D$ and right covariance matrix Ω , i.e.

$$(4.10) \quad [\mathcal{D}_s \mid \mathbf{B}, \mathbf{C}, \Sigma, \mathbf{Q}] \sim \mathcal{MN}_{n, jk}(\mathbf{H}_D \mathbf{B}, \mathbf{A}_D, \Omega)$$

Thus, using known ideas about the matrix normal distribution - see Dawid (1981), Carvalho and West (2007) - we write

$$(4.11) \quad [\mathcal{D}_s \mid \mathbf{B}, \mathbf{C}, \Sigma, \mathbf{Q}] = \frac{1}{(2\pi)^{\frac{njk}{2}} |\mathbf{A}_D|^{\frac{jk}{2}} |\Omega|^{\frac{n}{2}}} \exp \left\{ -\frac{1}{2} \text{tr} [\Omega^{-1} (\mathcal{D}_s - \mathbf{H}_D \mathbf{B})^T \mathbf{A}_D^{-1} (\mathcal{D}_s - \mathbf{H}_D \mathbf{B})] \right\}$$

The interpretation of the above is that the r -th row of $[\mathcal{D}_s \mid \mathbf{B}, \Sigma, \mathbf{C}, \mathbf{Q}]$ is multivariate normal with mean corresponding to row of the mean matrix $\mathbf{H}_D \mathbf{B}$ and with covariance matrix Ω . Rows r and ℓ

of $[\mathcal{D}_s | \mathbf{B}, \Sigma, \mathbf{C}, \mathbf{Q}]$ has covariance matrix $a(s_r, s_\ell) \Omega$. Similarly, the ℓ -th column of it is distributed as multivariate normal with mean being the ℓ -th column of $\mathbf{H}_D \mathbf{B}$ and with covariance matrix $\omega_{\ell, \ell} \mathbf{A}_D$, where $\omega_{r, \ell}$ denotes the (r, ℓ) -th element of Ω . The covariance between columns r and ℓ is given by the matrix $\omega_{r, \ell} \mathbf{A}_D$.

4.2. Predicting $s^{(new)}$. In order to predict the unknown Milky Way model parameter vector $s^{(new)}$ when the input is the measured real velocity vector $\mathbf{v}^{(test)}$, we write the posterior predictive distribution of $s^{(new)}$, given the real velocity matrix $\mathbf{v}^{(test)}$, the training data $\mathcal{D}_s$ and all the known targets, i.e. the chosen design vectors $s_1^*, \dots, s_n^*$. This posterior predictive is usually computed by integrating over all unknown model parameters. While it is possible to analytically integrate over $\mathbf{B}$ and $\mathbf{C}$, Σ and $\mathbf{Q}$ cannot be analytically integrated out. In fact, we find it useful to learn the d smoothing parameters (the diagonal elements of $\mathbf{Q}$), given the data. Thus, one useful advantage of our method is that the smoothness of the process does not need to be imposed by hand, but can be learnt from the data, if desired.

Given our learning of $s^{(new)}$, Σ and $\mathbf{Q}$, we rephrase our motivation as seeking to compute the joint posterior probability of $s^{(new)}$, $\mathbf{Q}$ and Σ , conditional on the real data and the training data, for a choice of the design matrix. In fact, we achieve a closed form expression of this joint posterior of $s^{(new)}$, $\mathbf{Q}$ and Σ , by integrating over the other hyperparameters, namely, the amplitude of the mean function ($\mathbf{B}$) and the matrix $\mathbf{C}$ that bears information about covariance among stars, for a given component of the velocity, at a fixed value of $\mathbf{S}$. From this closed form expression, the marginal posterior probability densities of $\mathbf{Q}$, Σ and any of the d components of the $s^{(new)}$ vector can be obtained, using the transformation based MCMC sampling method (Dutta and Bhattacharya, 2011) that we adopt.

For a given choice $s_1^*, \dots, s_n^*$ of the design vectors, the posterior distribution $[s^{(new)}, \Sigma, \mathbf{Q} | \mathbf{v}^{(test)}, \mathcal{D}_s]$ is sought, by marginalising the posterior $[s^{(test)}, \Sigma, \mathbf{Q}, \mathbf{B}, \mathbf{C} | \mathbf{v}^{(test)}, \mathcal{D}_s]$ over the process matrices $\mathbf{B}$ and $\mathbf{C}$.

4.3. Priors used. We use uniform prior on $\mathbf{B}$ and a simple non-informative prior on $\mathbf{C}$, namely, $\pi(\mathbf{C}) \propto |\mathbf{C}|^{-(j+1)/2}$. As for the priors on the other parameters, we assume uniform prior on $\mathbf{Q}$ and use the non-informative prior $\pi(\Sigma) \propto |\Sigma|^{-(k+1)/2}$. The prior information available in the astrophysical literature (Chakrabarty, 2007) on $s^{(new)}$ suggests uniform priors on all components of the $s^{(new)}$ vector (see Section 5.4 for greater details).

4.4. Posterior of $s^{(new)}$ given training and test data. Since our interest lies in the prediction of $s^{(new)}$, given the real test data and the simulated training data, as well as in learning the smoothness parameter matrix $\mathbf{Q}$ and the matrix Σ that bears the covariance for a given star, we compute the posterior distribution $[s^{(new)}, \mathbf{Q}, \Sigma | \mathbf{v}^{(test)}, \mathcal{D}_s]$. As expressed above, we achieve this by writing $[s^{(new)}, \mathbf{B}, \mathbf{C}, \mathbf{Q}, \Sigma | \mathbf{v}^{(test)}, \mathcal{D}_s]$ and marginalise over $\mathbf{B}$ and $\mathbf{C}$.

To construct an expression for this posterior distribution, we first collate the training and test data to construct the augmented data set $\mathcal{D}_{aug}^T = (\mathbf{v}_1^T; \dots; \mathbf{v}_n^T; (\mathbf{v}^{(test)})^T)$. Then the set of values of the model parameter vector $\mathbf{S}$ that supports $\mathcal{D}_{aug}$ is $\{s_1^*, \dots, s_n^*, s^{(new)}\}$ of which only $s^{(new)}$ is unknown. We define $\mathbf{H}_{\mathcal{D}_{aug}}^{((n+1) \times m)} := [\mathbf{h}^{(m \times 1)}(s_1^*), \dots, \mathbf{h}^{(m \times 1)}(s_n^*), \mathbf{h}^{(m \times 1)}(s^{(new)})]$, where our choice of the functional form of $\mathbf{h}(\cdot)$ has been given in Section 4 and we also set $m = d + 1$. Also, we define $\mathbf{A}_{\mathcal{D}_{aug}}^{((n+1) \times (n+1))} := [\exp\{-(s'_i - s'_{i'})^T \mathbf{Q} (s'_i - s'_{i'})\}]$ where s'_i and $s'_{i'}$ are members of the set $\{s_1^*, \dots, s_n^*, s^{(new)}\}$.

The priors used on $\mathbf{B}$, $\mathbf{C}$, $\mathbf{Q}$, Σ and $s^{(new)}$ are listed in Section 8. Using these, we observe that

$$\begin{aligned}
& [s^{(new)}, Q, B, \Sigma, C \mid v^{(test)}, \mathcal{D}_s] \propto [\mathcal{D}_{aug} \mid B, \Sigma, C, Q, s^{(new)}][B, \Sigma, C, Q, s^{(new)}] \\
& = |A_{\mathcal{D}_{aug}}|^{-\frac{jk}{2}} |\Sigma|^{-\frac{j(n+1)+k+1}{2}} |C|^{-\frac{k(n+1)+j+1}{2}} \\
& \times \exp \left[-\frac{1}{2} \text{tr} \left\{ \Omega^{-1} \mathcal{D}_{aug}^T M_{aug} \mathcal{D}_{aug} \right\} \right] \\
& \times \exp \left[-\frac{1}{2} \text{tr} \left\{ \Omega^{-1} \left\{ B - (H_{\mathcal{D}_{aug}}^T A_{\mathcal{D}_{aug}}^{-1} H_{\mathcal{D}_{aug}})^{-1} H_{\mathcal{D}_{aug}}^T A_{\mathcal{D}_{aug}}^{-1} \mathcal{D}_{aug} \right\}^T \right. \right. \\
& \quad \left. \left. H_{\mathcal{D}_{aug}}^T A_{\mathcal{D}_{aug}}^{-1} H_{\mathcal{D}_{aug}} \left\{ B - (H_{\mathcal{D}_{aug}}^T A_{\mathcal{D}_{aug}}^{-1} H_{\mathcal{D}_{aug}})^{-1} H_{\mathcal{D}_{aug}}^T A_{\mathcal{D}_{aug}}^{-1} \mathcal{D}_{aug} \right\} \right\} \right]
\end{aligned} \tag{4.12}$$

where we have recalled Equation 4.11. As in the definitions of $H_{\mathcal{D}_{aug}}$ and $A_{\mathcal{D}_{aug}}$, the matrix M_{aug} is defined at the given choice of the design matrix, augmented by the unknown Milky Way model parameter.

Marginalising $[s^{(new)}, Q, B, \Sigma, C \mid v^{(test)}, \mathcal{D}_s]$ (as expressed in Equation 4.12) over B and C yields the joint posterior $[s^{(new)}, Q, \Sigma \mid v^{(test)}, \mathcal{D}_s]$, the form of which is obtained as follows:

$$\begin{aligned}
& [s^{(new)}, Q, \Sigma \mid v^{(test)}, \mathcal{D}_s] \\
& = \int \int [s^{(new)}, Q, B, \Sigma, C \mid v^{(test)}, \mathcal{D}_s] dB dC \\
& \propto |A_{\mathcal{D}_{aug}}|^{-\frac{jk}{2}} \{H_{\mathcal{D}_{aug}}\}^T \{A_{\mathcal{D}_{aug}}\}^{-1} \{H_{\mathcal{D}_{aug}}\}^{-\frac{jk}{2}} \times |\Sigma|^{-\frac{j(n+1-m)+k+1}{2}} |(n+1-m)k \hat{C}_{GLS,aug}|^{-\frac{(n+1-m)k}{2}}
\end{aligned} \tag{4.13}$$

We thus obtain a closed-form expression of the joint posterior of $(s^{(new)}, Q, \Sigma)$, given training and test data, for a given choice of the design matrix (Equation 4.13), up to a normalising constant. The $\mathcal{GP}$ prior is strengthened by the n number of samples taken from it at the training stage. We sample from the achieved posterior using MCMC techniques to achieve the marginal posterior probabilities of Q , Σ or any component of $s^{(new)}$, given all data. In fact, we implement the transformation-based MCMC method recently by Dutta and Bhattacharya (2011).

4.5. Errors in measurement. In our application, the errors in the measurement of the stellar velocities are small and will be ignored for the rest of the analysis. In general, when errors in the measurements that comprise the training data and the test data are not negligible, we assume Gaussian stellar velocity errors ε_t , in v_t , with $t = 1, 2, \dots$, such that $\varepsilon_t \sim \mathcal{N}_{jk}(\mathbf{0}, \varsigma)$, where $\varsigma = \Sigma_1 \otimes \Sigma_2$; Σ_1, Σ_2 being positive definite matrices. If both Σ_1 and Σ_2 are chosen to be diagonal matrices, then ς is a diagonal matrix; assuming same diagonal elements would simplify ς to be of the form $\varphi \times \mathbf{I}$, where $\mathbf{I}$ is the $jk \times jk$ -th order identity matrix. This error variance matrix ς must be added to Ω before proceeding to the subsequent calculations. TMCMC can then be used to update ς .

Our fully Bayesian approach quantifies the prior information in terms of the Gaussian process and coherently processes the information contained in the prior and the data to yield results that are in agreement with those obtained by previous approaches using very large datasets. Moreover, unlike previous approaches, our approach allows for neat probabilistic conclusions *a posteriori*.

5. Analysis of real data. In our application, $\mathbf{S}$ is a 2-dimensional vector, i.e. $d=2$. Also, $\mathbf{P}$ is a two-dimensional vector, i.e. $k=2$. We consider 50 simulated/real stellar vectors at any value of $\mathbf{S}$ i.e. $j=50$. Lastly, we work with n design vectors where $n = 216$.

5.1. Posterior inference using TMCMC. Motivated by the fact that the performance of traditional MCMC methods - including the Metropolis-Hastings algorithm - can be less than satisfactory in high dimensions, both in terms of convergence and computational time, [Dutta and Bhattacharya \(2011\)](#) proposed a novel methodology that works by constructing proposals that are deterministic bijective transformations of some arbitrary random vector drawn from a chosen distribution. [Dutta and Bhattacharya \(2011\)](#) show that for additive transformations, the TMCMC-based acceptance rate decreases at a slower rate compared to block random walk Metropolis algorithms. Furthermore, TMCMC includes the hybrid Monte Carlo (HMC) algorithm as a special case and in one-dimensional situations while it boils down to the Metropolis-Hastings algorithm with a specialised proposal mechanism.

In TMCMC, the random vector of which a proposal density is a transformation of, can be chosen to be of dimensionality between 1 and the dimensionality of the parameters under the target posterior. High-dimensional parameter spaces are explored by constructing bijective deterministic transformations of a low-dimensional random vector. Interestingly, whatever the transformation, the acceptance ratio in TMCMC does not depend upon the distribution of the chosen random vector. For our purpose, we shall consider TMCMC based on additive transformations, since [Dutta and Bhattacharya \(2011\)](#) show that these transformations require far less number of “move types” compared to non-additive transformations.

5.2. TMCMC algorithm. TMCMC allows updating the entire block $(\mathbf{s}^{(new)}, \mathbf{Q}, \mathbf{\Sigma})$ at the same time. The algorithm is as follows.

- (i) Initialise the unknown quantities by fixing arbitrarily initial values $(\mathbf{s}^{(new,0)}, \mathbf{Q}^{(0)}, \mathbf{\Sigma}^{(0)})$. In our case, $\mathbf{s}^{(new,0)} = (s_1^{(new,0)}, \dots, s_d^{(new,0)})$, $\mathbf{Q}^{(0)}$ is characterised by the initial values of the d smoothness parameters, which we denote by $\mathbf{b} := (b_1^{(0)}, \dots, b_d^{(0)})^T$ and $\mathbf{\Sigma}^{(0)}$ denotes the initial choice of the $k \times k$ matrix $\mathbf{\Sigma}$. $\mathbf{\Sigma}$ is decomposed into $\mathbf{L}\mathbf{L}^T$, where $\mathbf{L}$ is the appropriate lower-triangular matrix
- (ii) Let $\boldsymbol{\varphi} = ((\mathbf{s}^{(new)})^T, \mathbf{b}^T, (\mathbf{L}^*)^T)^T$, where $\mathbf{L}^*$ denotes the column vector consisting of the non-zero elements of $\mathbf{L}$.
- (iii) Next we propose $\epsilon \sim g(\cdot)I_{\{\epsilon > 0\}}$, where $g(\cdot)$ is some arbitrary distribution, and I denotes the indicator function. In our applications, we shall choose $g(\cdot) = N(0, 1)$, so that, $\epsilon > 0$ is drawn from a truncated normal distribution.
- (iv) Assume that at iteration t , the state of the unknown parameters is $(\mathbf{s}^{(new,t)}, \mathbf{Q}^{(t)}, \mathbf{\Sigma}^{(t)}) := \boldsymbol{\varphi}^{(t)}$. Update $\boldsymbol{\varphi}^{(t)}$ by setting, with probabilities π_j and $(1 - \pi_j)$, $\varphi_j^{(t+1)} = \varphi_j^{(t)} \pm c_j \epsilon$ (forward transformation) and $\varphi_j^{(t+1)} = \varphi_j^{(t)} - c_j \epsilon$ (backward transformation), respectively, where, for $j = 1, \dots, d$, π_j are appropriately chosen probabilities and c_j are appropriately chosen scaling factors. Assume that for $j_1 \in \mathcal{U}$, $\varphi_{j_1}^{(t)}$ gets the positive transformation, while for $j_2 \in \mathcal{U}^c$, $\varphi_{j_2}^{(t)}$ gets the backward transformation. Here $\mathcal{U} \cup \mathcal{U}^c = \{1, \dots, d^*\}$, where $d^* = 2d + \frac{k(k+1)}{2}$.

(v) We accept the new proposal $\varphi^{(t+1)}$ with acceptance probability

$$(5.1) \quad \alpha_\varphi = \min \left\{ 1, \frac{\prod_{j_1 \in \mathcal{U}} (1 - \pi_{j_1}) \prod_{j_2 \in \mathcal{U}^c} \pi_{j_2}}{\prod_{j_1 \in \mathcal{U}} \pi_{j_1} \prod_{j_2 \in \mathcal{U}^c} (1 - \pi_{j_2})} \times r_\varphi \right\}$$

where r_φ denotes the ratio of $\left[|\mathbf{A}_{D_{aug}}(\mathbf{s}_{aug})|^{-\frac{ik}{2}} |\mathbf{H}_{D_{aug}}^T(\mathbf{s}_{aug}) \mathbf{A}_{D_{aug}}^{-1}(\mathbf{s}_{aug}) \mathbf{H}_{D_{aug}}(\mathbf{s}_{aug})|^{-\frac{ik}{2}} \right] \times \left[|\Sigma|^{-\frac{k(n+1-m)+k+1}{2}} |(n+1-m)k \hat{\mathbf{C}}_{GLS,aug}|^{-\frac{(n+1-m)k}{2}} \right]$, evaluated at the new value ($\varphi^{(t+1)}$) and the current value ($\varphi^{(t)}$) of φ respectively. We only need to bear in mind that the acceptance probability is zero if $b_j \leq 0$ for any j or if any diagonal element of $\mathbf{L}$ is negative.

5.3. Details of our implementation of TMCMC. In our work we use the following proposal mechanism: at each iteration we generated $\epsilon \sim \mathcal{N}(0, 1)I_{\{\epsilon > 0\}}$. Then, with equal probabilities, for $\ell = 1, 2$, we set $s_\ell^{(new, t)} = s_\ell^{(new, t-1)} \pm \epsilon$ and $b_\ell^{(t)} = b_\ell^{(t-1)} \pm 0.007\epsilon$. For updating Σ , we decompose $\Sigma = \mathbf{L}\mathbf{L}'$, but here we set the scale factor to be 0.07 for all the non-zero elements, using the additive transformation with equal move-type probabilities.

Following Section 5.2 we update all the unknowns in a single block, using additive transformations of a one-dimensional random variable. This, along with adequate choice of the scale parameters, ensures fast computations and reasonable convergence.

For proper choices of the scale parameters of the additive transformation and the initial values of the parameters we conducted several initial ‘‘pilot’’ TMCMC runs of length around 100,000, starting with arbitrary initial values and guesses of the scale parameters such that all the runs converged to the same distribution as indicated by informal diagnostics such as trace plots. For the final TMCMC run, we chose those scale parameters that yielded the best convergence (with respect to empirical diagnostics such as trace plots) among the pilot runs, and selected the final values of the parameters obtained in this best pilot run as the initial values for the final run of TMCMC. The pilot runs yielded the proposal mechanism that we worked with.

Once the proposal mechanism and the initial values are thus decided, we then discarded the next 100,000 iterations of our final TMCMC run as burn-in and stored the next 1,000,000 iterations for inference. For each model it took approximately 6 hours on a laptop to generate 1,100,000 TMCMC iterations.

5.4. Results from real data. The prior information available in the astrophysical literature (Chakrabarty, 2007) for the radial distance of the Sun from the galactic centre, $R_\odot \in [1.7, 2.3]$ and the angular separation of the Sun-Galactic centre line from the chosen long-axis of the elongated bar, $\Phi_\odot \in [0^\circ, 90^\circ]$.

The radial coordinate values are expressed in units of the ‘‘co-rotation radius’’ R_{CR} of the central bar in the Milky Way disc. This is the radius at which the rotational frequency Ω_b of the rigidly rotating bar, equals the radius-dependent rotational frequency $\Omega(R)$ of the stars at distance R from the centre of the Galaxy, which in turn is determined by the choice of the model for the gravitational field of the Milky Way disc that is one of the inputs in all 4 astrophysical models, the dynamical simulations of which produce the 4 training data sets that we implement in our work. In the model for the gravitational field of the Milky Way disk that we work with, this rotational frequency $\Omega(R)$ of the stars is defined as $\frac{v_0}{R}$, where v_0 is a constant that is set to unity in the simulation models of Chakrabarty (2007) and Ω_b is set to unity; thus $R_{CR} = 1$ in this scaled simulation model. All spatial lengths in our work are expressed in units of R_{CR} .

Astrophysical literature (Binney and Tremaine, 1987) offers the prior information that the radial location of the Sun is around $1.7R_{CR}$, which corresponds to the location of another major resonance - the Outer Lindblad Resonance - between the bar rotating at frequency Ω_b and the frequency of stars in the model Galactic disk. In particular, the training data that we use herein, was obtained by Chakrabarty (2007), by choosing the solar radial location from the interval $[1.7, 2.3]$. This explains the motivation for the aforementioned bounds on $R_\odot$, used in our work.



To connect the used model to real astronomical units, we need to scale $\Omega(R)$ as in the simulation units, to the stellar rotational frequency, as measured in real astronomical units. This calls for the scaling of the constant v_0 to astronomical units, and of the galactocentric radial distance R , measured on this scale in which $R_{CR} = 1$, to astronomical units. The Sun is a star for which the galactocentric radial location can be learnt in units of R_{CR} using our inverse Bayesian learning strategy presented above. Now, the constant v_0 is known in real units as 220 km s^{-1} (Binney and Merrifield, 1998). Independent astronomical estimates of the galactocentric solar radial location have led to the standard value of 8 kiloparsecs (Binney and Merrifield, 1998), where kiloparsec is an astronomical unit of distance. Then all distances encountered in the used model is connected to real astronomical unit of kiloparsec by scaling by the factor of $8/\hat{R}_\odot$ kiloparsec, where $\hat{R}_\odot$ is the solar radial location that we learn here. The co-rotation radius is then scaled from the simulation units to real units as $1 \times 8/\hat{R}_\odot$ kiloparsecs. Then from the definition of co-rotation, the bar rotational frequency Ω_b , scaled to real units, stands at $220/(8/\hat{R}_\odot) \text{ km s}^{-1} \text{ kpc}^{-1}$. Learning the rotational frequency of the bar is the ulterior benefit of learning the solar radial location as in our approach.

The azimuthal separation between the long axis of the bar and the line that joins the Sun to the Galactic centre, is also suggested in past astronomical modelling work to be an acute angle (Chakrabarty, 2007; Englmaier and Gerhard, 1999; Fux, 2001). Indeed, the training data used here was generated in simulations performed by Chakrabarty (2007), in which $\Phi_\odot$ is chosen from the interval $[0, 90^\circ]$. This motivates the consideration of the interval of $[0, 90^\circ]$ for the azimuthal location of the Sun.

Given the bounds on $R_\odot$ and $\Phi_\odot$ presented above, in our TMCMC algorithm, we reject those moves that suggest $R_\odot$ and $\Phi_\odot$ values that fall outside these presented intervals.

The 4 astrophysical models of the Galaxy that we discuss here are marked by the same choice of the value of Ω_b and the background Galactic model parameters, while they are distinguished by the varying choices of the ratio $\Omega_s : \Omega_b$, where the Galactic spiral pattern rotates with frequency Ω_s . In fact, the astrophysical model *bar6* is the only one that does not include the influence of the spiral pattern while the other three astrophysical models include the influence of both the bar and the spiral. In fact, the astrophysical models *sp3bar3_18*, *sp3bar3* and *sp3bar3_25* suggest $\Omega_s : \Omega_b$ values of $18\Omega_b/55$, $22\Omega_b/55$, $25\Omega_b/55$. The physical effect of this choice is to induce varying levels of chaoticity in the 4 astrophysical models. Thus, Chakrabarty and Sideris (2008) confirmed that of the 4 models, *bar6* manifests very low chaoticity while *sp3bar3* manifests maximal chaos at high energies, though both *sp3bar3_18*, *sp3bar3_25* are comparably chaotic.

Ancillary real data needs to be brought in to judge the relative fit amongst the astrophysical base models. In fact, Chakrabarty (2007) brought in extra information to perform model selection. Such information was about the observed variance of the radial and azimuthal components of stellar velocities and this was used to rule out the model *bar6* as physically viable, though the other three models were all acceptable from the point of view of all such ancillary observations that are available. This led to the inference that $\Omega_s \in [18\Omega_b/55, 25\Omega_b/55]$. It is to be noted that the 4 data sets simulated at the 4 distinct astrophysical models, cannot be meaningfully combined since the data sets are based on mutually

FIG 1. Posteriors of $R_{\odot}$ in units of R_{CR} and $\Phi_{\odot}$ (in degrees) for the model *bar6*.FIG 2. Posteriors of $R_{\odot}$ in units of R_{CR} and $\Phi_{\odot}$ (in degrees) for the model *sp3bar3*.

contradicting base models of the system. Hence we enumerate the results from each of the 4 models individually.

The marginal posterior densities of $(R_{\odot}, \Phi_{\odot})$ corresponding to the 4 astrophysical models of the Milky Way, are shown in Figures 1, 2, 3 and 4. Table 1 presents the posterior mode, the 95% highest posterior density (HPD) credible region of $R_{\odot}$ and $\Phi_{\odot}$ respectively, associated with the four models. Here $R_{\odot}$ is expressed in the model units of length, i.e. in units of R_{CR} . $\Phi_{\odot}$ is expressed in degrees. The HPDs are computed using the methodology discussed in [Carlin and Louis \(1996\)](#). Disjoint HPD regions, characterise the highly multi-modal posterior distributions of the unknown location. Using the 95% HPDs of the learnt $R_{\odot}$ expressed in model units, and using the independently known astronomical measurement of the solar radial location as 8 kiloparsecs, the bar rotational frequency Ω_b is computed (see explanation above) in Table 1.

Summaries of the posteriors (mean, variance and 95% credible interval) of the smoothness parameters b_1, b_2 and Σ are presented in Tables 2, 3. Notable in all these tables are the small posterior variances



FIG 3. Posteriors of $R_{\odot}$ in units of R_{CR} and $\Phi_{\odot}$ (in degrees) for the model *sp3bar3_18*.FIG 4. Posteriors of $R_{\odot}$ in units of R_{CR} and $\Phi_{\odot}$ (in degrees) for the model *sp3bar3_25*.

TABLE 1

Summary of the posterior distributions of the radial component $R_{\odot}$ and azimuthal component $\Phi_{\odot}$ of the unknown observer location vector for the 4 base astrophysical models and the unknown bar rotational frequency Ω_b computed using the 95% HPDs on the learnt radial location $R_{\odot}$ in these models.

Model	$R_{\odot}$ (in units of R_{CR})		Ω_b (in $\text{kms}^{-1}/\text{kiloparsec}$)	$\Phi_{\odot}$	
	Mode	95% HPD	95% HPD	Mode	95% HPD
<i>bar6</i>	2.20	[2.04, 2.30]	[56.1, 63.25]	23.50	[21.20, 25.80]
<i>sp3bar3</i>	1.73	[1.70, 2.26] $\cup$ [2.27, 2.28]	[46.75, 62.15] $\cup$ [62.45, 62.7]	18.8	[9.6, 61.5]
<i>sp3bar3_18</i>	1.76	[1.70, 2.29]	[46.75, 62.98]	32.5	[17.60, 79.90]
<i>sp3bar3_25</i>	1.95	[1.70, 2.15]	[46.75, 59.12]	37.6	[28.80, 40.40]

TABLE 2
Summary of the posterior distributions of the smoothness parameters b_1, b_2 for the 4 models.

Model	b_1			b_2		
	Mean	Var	95% CI	Mean	Var	95% CI
<i>bar6</i>	0.9598155	3.15×10^{-9}	[0.959703, 0.959879]	1.005078	2.85×10^{-9}	[1.004985, 1.005142]
<i>sp3bar3</i>	0.8739616	6.72×10^{-7}	[0.872347, 0.875052]	1.003729	8.98×10^{-7}	[1.002500, 1.005500]
<i>sp3bar3_18</i>	0.9410686	1.46×10^{-5}	[0.938852, 0.955264]	0.999010	4.08×10^{-6}	[0.997219, 1.004945]
<i>sp3bar3_25</i>	0.7597931	5.64×10^{-10}	[0.759743, 0.759833]	0.992174	2.89×10^{-9}	[0.992067, 0.992246]

TABLE 3
Summary of the posterior distribution of the diagonal and one non-diagonal element of Σ , from the 4 base astrophysical models.

Model	σ_{11}	σ_{22}	σ_{12}
	95% CI	95% CI	95% CI
<i>bar6</i>	$[5.40 \times 10^{-5}, 4.0 \times 10^{-4}]$	$[6.20 \times 10^{-5}, 4.76 \times 10^{-4}]$	$[0, 1.30 \times 10^{-5}]$
<i>sp3bar3</i>	$[3.66 \times 10^{-3}, 1.03 \times 10^{-2}]$	$[6.53 \times 10^{-3}, 1.83 \times 10^{-2}]$	$[-6.40 \times 10^{-5}, 2.68 \times 10^{-4}]$
<i>sp3bar3_18</i>	$[1.45 \times 10^{-3}, 1.68 \times 10^{-1}]$	$[1.29 \times 10^{-3}, 1.50 \times 10^{-1}]$	$[-1.19 \times 10^{-4}, 2.16 \times 10^{-3}]$
<i>sp3bar3_25</i>	$[1.21 \times 10^{-4}, 5.69 \times 10^{-4}]$	$[1.13 \times 10^{-4}, 5.21 \times 10^{-4}]$	$[-1.00 \times 10^{-6}, 1.50 \times 10^{-5}]$

of the quantities in question; this is indicative of the fact that the data sets we used, in spite of the relatively smaller size compared to the astronomically large data sets used in the previous approaches in the literature, are very much informative, given our vector-variate $\mathcal{GP}$ -based Bayesian approach.

As discussed in Section 5.6, thanks to our Gaussian Process approach, the posterior of Σ should be close to the null matrix *a posteriori* if the choice of the design set and the number of design points are adequate. Quite encouragingly, Table 3, shows that indeed Σ is close to the null matrix *a posteriori*, for all the four models, signifying that the unknown velocity function has been learned well in all the cases.

5.5. Comparison with results in astrophysical literature. The posterior summaries that we have presented in the last section, for each of the 4 astrophysical models of the Milky Way, compare favourably with results obtained by Chakrabarty (2007), using the calibration method. However, the all important difference in our implementation is the vastly smaller data set that we needed to invoke, in order to achieve the learning of the two-dimensional vector $\mathcal{S}$ - in fact while in the calibration approach, the required sample size is of the order of 3,500, in our work, this number is 50. Thus, data sufficiency issues, when a concern, are well tackled by our method.

Upon the analyses of the viable astrophysical models of the Galaxy, Chakrabarty (2007) reported the result that $R_\odot \in [1.9375, 2.21]R_{CR}$ while $\Phi_\odot \in [0^\circ, 30^\circ]$, where these ranges correspond to the presented uncertainties on the estimates, which were however, rather unsatisfactorily achieved (see Section 4). The values of the components of $\mathcal{S}$, learnt in our work, overlap well with these results. As mentioned in Section 1, learning $R_\odot$ and $\Phi_\odot$ have crucial astronomical significance for the azimuthal separation of the Sun from the long axis of the bar and for the rotational frequency Ω_b of the bar in the Milky Way. That the models *sp3bar3_18*, *sp3bar3* and *sp3bar3_25* are distinguished by distinct values of the ratios of the rotational frequencies of the spiral pattern to the bar in the Galaxy, the derived estimate for Ω_b (Table 1), in turn suggests possible credible regions on the value of the frequency Ω_s of the Milky Way spiral.

However, the concurrence of our results with the results reported in astrophysical literature goes beyond just the summaries of the posteriors of the solar position vector; remarkable correlation can be no-

ticed between the measure of chaos in these 4 astrophysical models - as estimated by (Chakrabarty and Sideris, 2008) - and the multi-modality of the posterior distribution of $\mathbf{S}$ that we advance. Chakrabarty and Sideris (2008) suggest minimum chaos in the *bar6* model compared to the other three, while we notice the posteriors of both $R_\odot$ and $\Phi_\odot$ in this model to be the unimodal - in fact the posteriors of $R_\odot$ and $\Phi_\odot$ are unimodal only for this model, out of the 4 astrophysical models that we use to illustrate the efficacy of our method. Perhaps more importantly, the *sp3bar3* model is noticed to manifest maximum (even global) chaoticity, on theoretical grounds by Chakrabarty (2007), backed by the chaos quantification at higher energies (Chakrabarty and Sideris, 2008). Likewise, in our work, the posterior distributions for $R_\odot$ and $\Phi_\odot$ are most multi-modal in this model, compared to the other three. The models *sp3bar3.18* and *sp3bar3.25* are considered to be of intermediate chaoticity and we find these to correspond to posterior distributions (of $\mathbf{s}^{(new)}$) that are multi-modal, though less so, than that for the model *sp3bar3*.

The equivalent in the work by Chakrabarty (2007), of our marginal posterior distributions of $(R_\odot, \Phi_\odot)^T$ is the multi-modality of the distribution of the p -value of the chosen test statistic in the space of $\mathbf{S}$, as used by Chakrabarty (2007) to test for the null that the observed velocity data is drawn from the density in the velocity space, estimated from the i -th simulated velocity data, for $i = 1, \dots, n$. When this distribution of the p -value was sought in the space of $\mathbf{S}$, the distribution was indeed found to be most multi-modal for the *sp3bar3* case while the modes were found to be less scattered than for the other three models. In line with this result, in our work we notice that the posterior probability of $\mathbf{S}$ is most multi-modal given the training data obtained with astrophysical model *sp3bar3*. The exact physical reason for the correlation between chaos in the astrophysical models and the multi-modality of the posterior distribution of $\mathbf{S}$ can be qualitatively understood if we begin with the premise that increased chaos gives rise to a more clustered distribution of the 2-dimensional stellar velocity vectors. This in turn implies that to an observer seated at any location on the Milky Way disk, the distribution of the velocities of n stars appears to be more multi-modal in nature than when there was less chaos (in the Galactic phase space). In other words, the distribution of the jk -dimensional $\mathbf{v}_i$ vectors is more multi-modal when the training data set is constructed using the *sp3bar3* model than the *bar_6* one. Consequently, the multi-modality in the achieved posterior probability density of $\mathbf{s}^{(new)}$ is higher in the former case than in the latter.

Another point that merits mentions is that the estimates of $R_\odot$ and $\Phi_\odot$ presented by Chakrabarty (2007) exclude the model *sp3bar3* which could not be used to yield estimates given the highly scattered nature of the corresponding p -value distribution. Likewise, in our work, the same model manifests maximal multi-modality amongst the others, but importantly, our approach allows for the representation of the full posterior density using which, the computation of the 95% HPDs is performed.

5.6. Issue related to data sufficiency. It was noted in Section 5.4 above that in spite of our relatively smaller data sets, our learning of the unknown location vector is in accordance with the relevant results obtained by other approaches existing in the astrophysics literature. In this connection it is important to clarify how inference of the unknown quantities might be affected by the dimensionality of $\xi(\cdot)$ which is in turn set by the number of stars chosen at any $\mathbf{s}_i$, $i = 1, \dots, n$. Note that, irrespective of dimensionality, the Bayesian-ly predicted function must pass through all the points of the training data set. It is the Σ matrix that bears information about covariance for a given star, (at a given value of the model parameter vector $\mathbf{S}$), in our model. That the elements of the learnt Σ matrix are learnt to be small, and our learnt $\mathbf{s}^{(new)}$ are astrophysically interpretable, corroborate the stability of our inversion given the n repeated samples. Such instabilities in inversion (Banerjee et al., 2008) can become effective for the following reason. If a particular functional value is supported by some argument, inclusion of more

(different) functions can change this value such that now it supports the old functions as well as the new functions. It is because we seek verification under such a situation that it becomes useful to learn the covariance matrix Σ in the data (while the column covariance matrix C can be analytically integrated over). In fact, marginalisation over C gives rise to a parameter space that is more informed by n than j , i.e. more informed by the number of known values of S at which data is simulated than the number of stars used at a given value of S .

Also, each element of $\xi(\cdot)$ has the same smoothness parameters for two given values of the Milky Way model parameter vector (see Equation 4.9). Thus, increasing the number of stars while maintaining the number of design points a constant, will fail to improve inference about the smoothness parameters. We realise this by recalling that since the smoothness parameters determine the smoothness of the function $\xi(\cdot)$ with respect to the sample path - design points being the discretised version - information about the smoothness parameters is enhanced by increasing the number of design points rather than the number of stars.

The arguments presented above show that the aim of improving the quality of inference on Σ and Q , is better served, if we increase the number n of design points in the used training data set, than the number of stars at each choice of the value s_i of the model parameter, $i = 1, \dots, n$. Understanding these data-sufficiency issues turns out to be very rewarding from the computational point of view – increasing the number of stars implies increasing the dimension of the matrix C , which would have made our computation prohibitively slow, but keeping the dimension of C at some reasonable level, and increasing n is computationally much less expensive.

In our case, results for the unknown location are obtained by our approach with 50×2 velocity functions, which is a reasonable dimensionality, along with 216 well-chosen design points. Our results, which are astrophysically interpretable, demonstrate that these choices are adequate.

5.7. Benefits of the method. An important benefit of our work is that it allows for the learning of a high-dimensional model parameter vector. Till now, a rigorous statistical learning of the Milky Way model parameter vector, from the inversion of available - real and simulated - stellar velocity data, has not taken place. Additionally, all previous attempts have acknowledged the need for including other Milky Way model parameters in their respective modelling strategies, but the lack of rigour in the same did not allow for the learning exercise to be performed in the context of a high-dimensional model parameter vector. For example, it is of substantial astronomical value if one could learn strengths of the perturbations due to the Galactic features as well other structural parameters and parameters that determine their dynamical behaviour (such as the ratio $\Omega_s : \Omega_b$). In our method, extension to such a high-dimensional S is easily possible, as long as astrophysical simulations carried out at varying values of these parameters are available. In fact, it is the matrix $A_{\mathcal{D}_{aug}}(\cdot)$ discussed above, that will be affected by the increase in the value of d ; the computational complexity involved in the computation of this matrix scales linearly with d .

The implication of our demonstrated success with a smaller data set is profound. Firstly, our method is advanced as a template for the analysis of the stellar phase space data that is available for the Milky Way, with the aim of learning a high-dimensional Galactic parameter vector; by extending the scope of the dynamical simulations of the Galaxy, performed on different astrophysical models of the Milky Way, the Milky Way models will be better constrained. At the same time, the planned flight of the GAIA - a mission of the European Space Agency - later in 2012, is set to provide large sets of phase space data all over the Milky Way. Our method, in conjunction with astrophysical models, can allow for fast learning of local and global model parameters of the Galaxy. In fact, the greater strength of our method lies in

its proposed applicability to galaxies other than our own, for which, only small data sets ($\lesssim 100$) can be expected. The learning of global model parameters of such systems is possible with a method such as ours, once data generated from simulations of such systems are available.

6. Discussions. We have presented a new method to allow for inverse Bayesian learning of model parameter vectors, given real and simulated data, using a vector-variate Gaussian Process that was motivated as an alternative for a matrix-variate $\mathcal{GP}$ of corresponding dimensions. The biggest advantage of our method is that it can work even when data sufficiency might be a concern and when making inference about a high-dimensional model parameter vector is infeasible in other methods. The learning is rendered possible, owing to the matrix-variate Gaussian Process that we introduce here and successfully demonstrate an application of in an astronomical context.

We saw in Section 5.7 that computational complexity scales only linearly with the dimensionality of the unknown model parameter $\mathbf{S}$. Thus, porting a training data comprised of n independent values of $\mathbf{S}$, $\mathbf{s}_i$, $i = 1, \dots, n$, where $\mathbf{s}_i$ is a d -dimensional vector, $d > 2$, is not going to render the computational times infeasible. This allows for the learning of high-dimensional model parameter vectors in our method.

In contrast to the situation with increasing the dimensionality of the unknown model parameter, increasing the dimensionality of the measurable will but imply substantial increase in the run time, since the relevant computational complexity then scales non-linearly, as about $O(k^3)$, (in addition to the cost of k square roots), where k is the dimensionality of the measurable variable $\mathbf{P}$. This is because of the dimensionality of the aforementioned Σ matrix is $k \times k$, and the inverse of this enters the computation of the posterior. Thus, for example, increasing the measurable from a 2-dimensional to 4-dimensional vector increases the run time 8-fold, which is a large jump in the required run time. However, for most applications, we envisage the expansion of the dimensionality of the unknown model parameter, i.e. d , rather than that of the measurable, i.e. k . Thus, the method is expected to yield results within acceptable time frames, for most practical applications.

The other major benefit of our work is that it allows for organic learning of the smoothness parameters, rather than results being subject to ad hoc choices of the same.

Inverse learning similar to our application, is essentially of wide applicability - for example, one might need to learn the required dosage of a drug in order to achieve a cure efficacy that is judged optimal in curing a disease in a sample of j patients. This can be done using available data from a set of trials (the training data). Thus, in this scenario, n number of trials might be performed, each with j patients, such that changes in k characteristics of each patient are monitored where these measured “characteristics” reflect the efficacy of the administered dosage of the medication in affecting k different symptoms of the ailment. Thus, the efficacy parameter $\mathbf{E}$ is a k -dimensional vector and we have data on the value of $\mathbf{E}$ of j patients, from n different trials that are distinguished by the n values of the dosage vector $\mathbf{D}$. While dosage of the medication could be perceived as a scalar, it is also possible that it is rendered vector-valued, as can be the case, if there is reason to vary the dosage of the drug administered at different times of the day. In general, we consider $\mathbf{D}$ to be a d -dimensional vector. The test data is the data on the optimal improvement (cure efficacy) of k symptoms of these j patients when the administered dosage is the unknown $\mathbf{D}^{(test)}$. Using the $j \times k$ -dimensional efficacy matrices, attained for each of the n trials for the n values of the dosage vector, $\{\mathbf{D}_i\}_{i=1}^n$, $\mathbf{D}^{(test)}$ can be learnt. Here, the matrix-valued information on the continuous parameter $\mathbf{E}$ (cure efficacy) is perceived as an unknown function of the dosage parameter vector which too is continuous in nature. We can realise this unknown function as a realisation of a Gaussian Process, after the method that we present above.

Alternatively, the price of a commodity might need to be fixed, given a parametrisation of the current demand for the item, guided by training data comprising such a demand parameter and the price, at time points in the past.

In another astrophysical context, [Chakrabarty and Jackson \(2009\)](#) learnt the distribution of the density of the total (i.e. luminous+dark) gravitational matter $\rho : \mathbb{R}^3 \rightarrow \mathbb{R}$, $\rho(x, y, z) \geq 0$, in distant galaxies belonging to an ubiquitous astronomical class of real galactic systems that is parametrised by the shape parameter $n > 0$ and a length scale parameter $X_e \geq 0$. The measurements include the observed galactic image and central velocity dispersion measurement in the galaxy, while training data comprising synthetic image data and central velocity dispersion, generated from simulations of such systems carried out at chosen values of n , X_e and other relevant model variables. In such applications, as the measurables are rendered vectors and the data a matrix, inverse learning is possible using the method that we present here.

As more Galactic simulations spanning a greater range of model parameters become available, the rigorous learning of such Milky Way parameters using our method will become possible, given the available stellar velocity data. This will enhance the quality of our knowledge about our own galaxy. That our method allows for such learning even for under-abundant systems, is encouraging for application of a similar analysis to galaxies other than our own, in which system parameters may be learnt using the much smaller available velocity data sets, compared to the situation in our galaxy.

Supplementary material

Section S-1 discusses the details of the dynamical simulations that lead to the training data set used in our supervised learning of the Milky Way model parameters. The posteriors of the matrices $\mathbf{B}$ and $\mathbf{C}$ that characterise the Gaussian Process under consideration are presented in S-2. S-3 contains details of a proof that the matrix $(n-m)k\hat{C}_{GLS}$ is positive definite. In Section S-4, a discussion about the batchwise updating within the TMCMC procedure is included.

Acknowledgments. We thank Prof. Ayanendranath Basu, Prof. Smarajit Bose, Mr. Pronoy K. Mondal, and Mr. Sayan Patra for helpful discussions. That the matrix $\hat{C}_{GLS}$ is positive definite has been proved formally by Mr. Pronoy Mondal. DC acknowledges a Warwick Centre for Analytical Sciences Fellowship under the support of which the work was initiated.

SUPPLEMENTARY SECTION FOR “BAYESIAN NONPARAMETRIC ESTIMATION OF MILKY WAY MODEL PARAMETERS USING A NEW MATRIX-VARIATE GAUSSIAN PROCESS BASED METHOD”

Throughout, we refer to our main manuscript as CBB.

7. Details of dynamical simulations of astrophysical models. In [Chakrabarty \(2007\)](#), the modelling involves the following. A sample of phase space coordinates $\{\mathbf{w}_0\}$, is drawn from a chosen (to mimic real disc galaxies’) phase space density $g(\mathbf{w})$ at $t = 0$, and is evolved in a (chosen) parametric Galactic gravitational potential $\Psi : \mathbb{R}^3 \times \mathcal{T} \rightarrow \mathbb{R}$, where the time variable $T \in \mathcal{T} \subset \mathbb{R}_{\geq 0}$. $\Psi(\mathbf{X}, t)$ is chosen to emulate a realistic background Galactic potential $\Psi_0(\mathbf{x}) \in \mathbb{R}$, perturbed by the chosen parametric forms (rigidly rotating, quadrupolar) of the gravitational potential of the bar ($\Psi_b(\mathbf{x}) \in \mathbb{R}$) and the (logarithmic spiral) gravitational potential of the spiral pattern ($\Psi_s(\mathbf{x}) \in \mathbb{R}$) in the Milky Way. The perturbation strengths of these features ($\varepsilon_b : \mathcal{T} \rightarrow \mathbb{R}$ and $\varepsilon_s : \mathcal{T} \rightarrow \mathbb{R}$) are slowly increased, to saturation values over time $T_s \in \mathbb{R}$, $T_s > 0$ which is chosen such that $T_s \ll T_{sim}$, where $T_{sim} \in \mathbb{R}$ is the total simulation time. Thus, in the q^{th} astrophysical model, $\Psi^{(q)}(\mathbf{x}, t) = \Psi_0^{(q)}(\mathbf{x}) + \varepsilon_b(t)\Psi_b^{(q)}(\mathbf{x}, t) + \varepsilon_s(t)\Psi_s^{(q)}(\mathbf{x}, t)$, for each $q = 1, \dots, 4$. At $T = T_{sim}$, orbits are sampled in phase and recorded stroboscopically in the rotating frame of the bar at times when $(\Omega_b - \Omega_s)t=0$.

In order to sort the orbits by the value of the solar location in a galactocentric coordinate frame, chosen ranges of $R \in [1.7, 2.3]$ in simulation units and $\Phi \in [0, 90]$ in degrees, are used to define a regular, 2-D polar grid with $N_R \times N_\phi$ cells of radial and azimuthal widths of δ_R and δ_ϕ , $\delta_R, \delta_\phi \in \mathbb{R}$. The i^{th} cell in this grid represents the i^{th} value of the model parameter vector, $\mathbf{S}^{(i)}$, with $n = N_R \times N_\phi$, $i = 1, \dots, N_R \times N_\phi$. In order to find stars that end up in the simulations with spatial coordinates within the interval around $\mathbf{s}^{(i)}$ that defines the i^{th} -cell, we identify stars with radius $\in [s_1^{(i)}, s_1^{(i)} + \delta_1)$ and azimuth $\in [s_2^{(i)}, s_2^{(i)} + \delta_2)$; these stars comprise the simulated velocity information $\mathbf{V}_i$ in the i^{th} -cell. This is performed for each i , $i = 1, \dots, n$. The same 2-D polar grid is used for each of the Milky Way astrophysical models.

8. Posteriors of B and C given training data. Here, we discuss the achievement of the posteriors of the process matrices B and C , given the training data. To write the posteriors, we use uniform prior on B and a simple non-informative prior on C , namely, $\pi(C) \propto |C|^{-(j+1)/2}$.

Recalling that the distribution of the training data $\mathcal{D}_s$ is matrix normal, we write the posterior of B as

$$(8.1) \quad [B \mid \Sigma, C, Q, \mathcal{D}_s] \sim \mathcal{MN}_{m,jk}(\hat{B}_{GLS}, (H_D^T A_D^{-1} H_D^{-1}, \Omega),$$

where, $\hat{B}_{GLS} := (H_D^T A_D^{-1} H_D)^{-1} (H_D^T A_D^{-1} \mathcal{D}_s)$; note that $H_D \hat{B}_{GLS}$ can be interpreted as the generalised least square (GLS) solution to the equation that given the chosen design vectors, the mean matrix $H_D B$ is equal to $\mathcal{D}_s$.

To write the posterior of C , for a chosen design set, we first define

$$(8.2) \quad M_D := A_D^{-1} - A_D^{-1} H_D [H_D^T A_D^{-1} H_D]^{-1} H_D^T A_D^{-1},$$

and $(\mathcal{D}_s^T \mathbf{M}_D \mathcal{D}_s)^{(jk \times jk)} := [\mathbf{M}_{it}^*; i, t = 1, \dots, k]$, where $\mathbf{M}_{it}^*$ is a matrix with j rows and j columns. Given Σ , we define $m = d + 1$ and ψ_{it}^{-1} as the (i, t) -th element of Σ^{-1} , so that $(n - m)k\hat{\mathbf{C}}_{GLS} := \sum_{i=1}^k \sum_{t=1}^k \psi_{it}^{-1} \mathbf{M}_{ti}^*$. Then it can be shown that for this given choice of the design vectors, the posterior distribution $[C|\Sigma, \mathbf{Q}, \mathcal{D}_s]$ is inverse Wishart with parameters $(n - m)k\hat{\mathbf{C}}_{GLS}$ and $(n - m)k$ (degrees of freedom).

9. Positive definiteness of $(n - m)k\hat{\mathbf{C}}_{GLS}$. Now marginalisation of the posterior $[\mathbf{s}^{(test)}, \Sigma, \mathbf{Q}, \mathbf{B}, \mathbf{C}|\mathbf{v}^{(test)}, \mathcal{D}_s]$ are sensitive to the posterior $[C|\Sigma, \mathbf{Q}, \mathcal{D}_s]$. Thus, we need to ensure that $(n - m)k\hat{\mathbf{C}}_{GLS}$ is positive definite in order for the posterior of $\mathbf{C}$ to be well-defined. We now present the proof that $(n - m)k\hat{\mathbf{C}}_{GLS}$ is positive definite.

PROOF. Since $\mathcal{D}_s^T \mathbf{M}_D \mathcal{D}_s = [\mathbf{M}_{it}^*; i, t = 1, \dots, k]$ is positive definite, $\mathbf{B}^T \mathbf{M}_D \mathcal{D}_s = \mathbf{P} \mathbf{P}'$, where $\mathbf{P}$ is a lower triangular matrix with strictly positive diagonal entries. Writing $\mathbf{P} = (\mathbf{P}_1^T; \mathbf{P}_2^T; \dots; \mathbf{P}_k^T)^T$, where each $\mathbf{P}_t$ is of order $j \times j$, it then follows that $\mathbf{M}_{it}^* = \mathbf{P}_i' \mathbf{P}_t$. Hence, for any non-null j -component vector $\mathbf{s}$,

$$(9.1) \quad \sum_{i=1}^k \sum_{t=1}^k \sigma_{it}^{-1} \mathbf{s}^T \mathbf{M}_{ti}^* \mathbf{s} = \sum_{i=1}^k \sum_{t=1}^k \sigma_{it}^{-1} (\mathbf{P}_i \mathbf{s})^T (\mathbf{P}_t \mathbf{s}).$$

Here σ_{it} is the it -th element of the matrix $\sigma_D(\cdot) := [a(\cdot, \mathbf{s}_1), \dots, a(\cdot, \mathbf{s}_n)]^T$. The right hand side of the last equation is $\mathbf{y}^T (\Sigma^{-1} \otimes \mathbf{I}_j) \mathbf{y}$, where $\mathbf{y} := (\mathbf{y}_1^T, \dots, \mathbf{y}_k^T)^T$, with $\mathbf{y}_t = \mathbf{P}_t \mathbf{s}$, and $\mathbf{I}_j$ is the j -th order identity matrix. Since Kronecker product of positive definite matrices yields positive definite matrices, it follows that the last expression is positive, implying that $(n - m)k\hat{\mathbf{C}}_{GLS}$ is positive definite. $\square$

10. Batchwise TMCMC based on additive transformations. Using TMCMC we can update the parameters in batches as follows:

- (i) Initialise the unknown quantities by fixing arbitrarily initial values $(\mathbf{s}^{(test,0)}, \mathbf{Q}^{(0)}, \Sigma^{(0)})$. In our case, $\mathbf{s}^{(test,0)} = (s_1^{(test,0)}, \dots, s_d^{(test,0)})$, $\mathbf{Q}^{(0)}$ is characterised by the initial values of the d smoothness parameters, which we denote by $(b_1^{(0)}, \dots, b_d^{(0)})$ and $\Sigma^{(0)}$ denotes the initial choice of the $k \times k$ matrix Σ .
- (ii) Assume that at iteration t , the state of the unknown parameters is $(\mathbf{s}^{(test,t)}, \mathbf{Q}^{(t)}, \Sigma^{(t)})$.
- (iii) (a) Propose $\epsilon \sim g(\cdot) I_{\{\epsilon > 0\}}$, where $g(\cdot)$ is some arbitrary distribution, and I denotes the indicator function. In our applications, we shall choose $g(\cdot) = N(0, 1)$, so that, $\epsilon > 0$ is drawn from a truncated normal distribution. Update the d components of location $\mathbf{s}^{(test)}$ by setting, with probabilities π_j and $(1 - \pi_j)$, $s_j^{(test,t+1)} = s_j^{(test,t)} + c_j \epsilon$ (forward transformation) and $s_j^{(test,t+1)} = s_j^{(test,t)} - c_j \epsilon$ (backward transformation), respectively, where, for $j = 1, \dots, d$, π_j are appropriately chosen probabilities and c_j are appropriately chosen scaling factors. Assume that for $j_1 \in \mathcal{S}$, $\mathbf{s}_{j_1}^{(test,t)}$ gets the positive transformation, while for $j_2 \in \mathcal{S}^c$, $\mathbf{s}_{j_2}^{(test,t)}$ gets the backward transformation. Here $\mathcal{S} \cup \mathcal{S}^c = \{1, \dots, d\}$.
- (b) Accept the new proposal $\mathbf{s}^{(test,t+1)}$ with acceptance probability

$$(10.1) \quad \alpha_s = \min \left\{ 1, \frac{\prod_{j_1 \in \mathcal{S}} (1 - \pi_{j_1}) \prod_{j_2 \in \mathcal{S}^c} \pi_{j_2}}{\prod_{j_1 \in \mathcal{S}} \pi_{j_1} \prod_{j_2 \in \mathcal{S}^c} (1 - \pi_{j_2})} \times r_s \right\}$$

where r_s denotes the ratio of $|A_{D_{aug}}|^{-\frac{jk}{2}} |H'_{D_{aug}} A_{D_{aug}}^{-1} H_{D_{aug}}|^{-\frac{jk}{2}} \times |\Sigma|^{-\frac{k(n+1-m)+k+1}{2}} | (n+1-m)k \hat{C}_{GLS,aug} |^{-\frac{(n+1-m)k}{2}}$, evaluated at the new value $(s^{(test,t+1)})$ and the current value $(s^{(test,t)})$ of $s^{(test)}$ respectively, with b and Σ remaining fixed at the current values $b^{(t)}$ and $\Sigma^{(t)}$, respectively. In our applications, we shall choose $\pi_j = 1/2 \forall j$, which simplifies the acceptance ratio by eliminating the first factor altogether.

- (c) If the new proposal $s^{(test,t+1)}$ is not accepted, we set $s^{(test,t+1)} = s^{(test,t)}$.
- (iv) Using the same TMCMC strategy, update b . Note that the probability of acceptance is zero if some components of the proposed value of b is negative.
- (v) Update Σ by first decomposing it into LL' , where L is the appropriate lower-triangular matrix. Then update all the non-zero elements of L in a single block using the same TMCMC strategy. The form of the acceptance ratio remains same. Again, reject the moves that yield negative diagonal elements.
- (vi) Cycle over steps (ii)–(v) for $K_1 + K_2$ iterations, assuming that the Markov chain converges after K_1 iterations. Store the realisations $\left\{ \left(s^{(test,t)}, Q^{(t)}, \Sigma^{(t)} \right) : t = K_1 + 1, \dots, K_2 \right\}$ as samples obtained (approximately) from $[s^{(test)}, Q, \Sigma \mid \mathcal{D}_{aug}]$. In particular, the realisations $\{s^{(test,t)} : t = K_1 + 1, \dots, K_2\}$ are samples (approximately) from the marginal distribution $[s^{(test)} \mid V^{(test)}, \mathcal{D}_{aug}]$.

References.

- A. O'HAGAN, (1978). Curve Fitting and Optimal Design for Prediction. *Journal of the Royal Statistical Society B* **40** 1-42.
- ANTOJA, T., VALENZUELA, O., PICHARDO, B., MORENO, E., FIGUERAS, F. and FERNÁNDEZ, D. (2009). Stellar Kinematic Constraints on Galactic Structure Models Revisited: Bar and Spiral Arm Resonances. *Astrophysical J. Letters* **700** L78-L82.
- BANERJEE, S., GELFAND, A. E., FINLEY, A. and SANG, H. (2008). Gaussian predictive process models for large spatial data sets. *Journal of the Royal Statistical Society, Series B* **70** 825-848.
- BINNEY, J. and MERRIFIELD, M. (1998). *Galactic Astronomy*. Princeton University Press, Princeton.
- BINNEY, J. and TREMAINE, S. (1987). *Galactic Dynamics*. Princeton University Press, Princeton.
- BLIGHT, B. J. N. and OTT, L. (1975). A Bayesian Approach to Model Inadequacy for Polynomial Regression. *Biometrika* **62** 79-88.
- CARLIN, B. P. and LOUIS, T. A. (1996). *Bayes and Empirical Bayes Methods for Data Analysis*. Chapman and Hall, London. Second Edition.
- CARVALHO, C. M. and WEST, M. (2007). Dynamic Matrix-Variate Graphical Models. *Bayesian Analysis* **2** 69-98.
- CHAKRABARTY, D. (2007). Phase Space around the Solar Neighbourhood. *Astronomy & Astrophysics* **467** 145.
- CHAKRABARTY, D. (2011). Galactic Phase Spaces. In *Proceedings of the 7th Workshop on Data Analysis in Astronomy—Science: IMAGE IN ACTION* Ettore Majorana Foundation and Centre for Scientific Culture, Erice, April 15-22, 2011.
- CHAKRABARTY, D. and JACKSON, B. (2009). Total Mass Distributions of Sersic Galaxies from Photometry & Central Velocity Dispersion. *Astronomy & Astrophysics* **498** 615.
- CHAKRABARTY, D. and SIDERIS, I. (2008). Chaos in Models of the Solar Neighbourhood. *Astronomy & Astrophysics* **488** 161.
- CRESSIE, N. A. C. (1993). *Statistics for Spatial Data*. Wiley, New York.
- DAWID, A. P. (1981). Some Matrix-Variate Distribution Theory: Notational Considerations and a Bayesian Application. *Biometrika* **68** 265-274.
- DEHNEN, W. (2000). The Effect of the Outer Lindblad Resonance of the Galactic Bar on the Local Stellar Velocity Distribution. *Astronomical Journal* **11** 800.
- DUTTA, S. and BHATTACHARYA, S. (2011). Markov Chain Monte Carlo Based on Deterministic Transformations. Submitted, available at <http://arxiv.org/abs/1106.5850>.
- ENGLMAIER, P. and GERHARD, O. (1999). Gas dynamics and large-scale morphology of the Milky Way galaxy. *Monthly Notices of the Royal Astronomical Society* **304** 512-534.
- FUX, R. (2001). Order and chaos in the local disc stellar kinematics induced by the Galactic bar. *Astronomy & Astrophysics* **373** 511-535.
- G. WALKER AND J. FORD, (1969). Amplitude Instability and Ergodic Behavior for Conservative Nonlinear Oscillator Systems. *Physical Review* **188** 416-432.
- JOHNSTON, K., BULLOCK, J. S. and STRAUSS, M. (2009). The Milky Way and Local Volume as Rosetta Stones in Galaxy Formation. In *astro2010: The Astronomy and Astrophysics Decadal Survey* **2010** 142.
- KUCINSKAS, A., LINDEGREN, L. and VANSEVICIUS, V. (2005). Beyond the Galaxy with Gaia: Evolutionary Histories of Galaxies in the Local Group. In *The Three-Dimensional Universe with Gaia* (C. TURON, K. S. O'FLAHERTY and M. A. C. PERRYMAN, eds.). *ESA Special Publication* **576**.
- LINDEGREN, L., BABUSIAUX, C., BAILER-JONES, C., BASTIAN, U., BROWN, A. G. A., CROPPER, M., HG, E., JORDI, C., KATZ, D., VAN LEEUWEN, F., LURI, X., MIGNARD, F., DE BRUIJNE, J. H. J. and PRUSTI, T. (2007). The Gaia mission: science, organization and present status. In *A Giant Step: from Milli- to Micro-arcsecond Astrometry, Proceedings IAU Symposium No. 248* (W. J. JIN, I. PLATAIS and M. A. C. PERRYMAN, eds.) 217-223.
- MATERN, B. (1986). *Spatial Variation (2nd ed.)*. Springer-Verlag, Springer.
- MINCHEV, I., QUILLEN, A. C., WILLIAMS, M., FREEMAN, K. C., NORDHAUS, J., SIEBERT, A. and BIENAYMÉ, O. (2009). Is the Milky Way ringing? The hunt for high-velocity streams. *Monthly Notices of the Royal Astronomical Society* **396** L56-L60.
- MINCHEV, I., BOILY, C., SIEBERT, A. and BIENAYME, O. (2010). Low-velocity streams in the solar neighbourhood caused by the Galactic bar. *Monthly Notices of the Royal Astronomical Society* **407** 2122-2130.
- NEAL, R. M. (1998). Regression and classification using Gaussian process priors (with discussion). In *Bayesian Statistics 6* (J. M. B. ET. AL, ed.) 475-501. Oxford University Press.
- RASMUSSEN, C. E. and WILLIAMS, C. K. I. (2006). *Gaussian Processes for Machine Learning*. The MIT Press, MIT.
- SANTNER, T. J., WILLIAMS, B. J. and NOTZ, W. I. (2003). *The design and analysis of computer experiments*. Springer Series in Statistics. Springer-Verlag, New York, Inc.

- SCHOLKOPF, B. and SMOLA, A. J. (2002). *Learning with Kernels*. MIT Press, MIT.
- SIMONE, R. D., WU, X. and TREMAINE, S. (2004). The stellar velocity distribution in the solar neighbourhood. *Monthly Notices of the Royal Astronomical Society* **350** 627.
- SNELSON, E. L. (2007). Flexible and efficient Gaussian process models for machine learning Doctoral Thesis, University of London.
- TILMANN GNEITING AND WILLIAM KLEIBER AND MARTIN SCHLATHER, (2010). Matern Cross-Covariance Functions for Multivariate Random Fields. *Journal of the American Statistical Association* **105** 1167-1177.